



SCHOOL OF ECONOMICS AND MANAGEMENT
FACULTY OF SOCIAL SCIENCES
AARHUS UNIVERSITY



CREATES
Center for Research in Econometric
Analysis of Time Series

CREATES Research Paper 2009-52

Jump-Robust Volatility Estimation using Nearest Neighbor Truncation

Torben G. Andersen, Dobrislav Dobrev and Ernst Schaumburg

School of Economics and Management
Aarhus University
Bartholins Allé 10, Building 1322, DK-8000 Aarhus C
Denmark

Jump-Robust Volatility Estimation using Nearest Neighbor Truncation

Torben G. Andersen, Dobrislav Dobrev, Ernst Schaumburg *

This version: October 31, 2009

Abstract

We propose two new jump-robust estimators of integrated variance based on high-frequency return observations. These MinRV and MedRV estimators provide an attractive alternative to the prevailing bipower and multipower variation measures. Specifically, the MedRV estimator has better theoretical efficiency properties than the tripower variation measure and displays better finite-sample robustness to both jumps and the occurrence of “zero” returns in the sample. Unlike the bipower variation measure, the new estimators allow for the development of an asymptotic limit theory in the presence of jumps. Finally, they retain the local nature associated with the low order multipower variation measures. This proves essential for alleviating finite sample biases arising from the pronounced intraday volatility pattern which afflict alternative jump-robust estimators based on longer blocks of returns. An empirical investigation of the Dow Jones 30 stocks and an extensive simulation study corroborate the robustness and efficiency properties of the new estimators.

JEL classification: C14; C15; C22; C80; G10.

Keywords: High-frequency data; Integrated variance; Finite activity jumps; Realized volatility; Jump robustness; Nearest neighbor truncation;

*Torben G. Andersen: Northwestern Univ.; NBER; CREATES; t-andersen@northwestern.edu
Dobrislav Dobrev: Federal Reserve Board of Governors, Dobrislav.P.Dobrev@frb.gov
Ernst Schaumburg: Federal Reserve Bank of New York, Ernst.Schaumburg@ny.frb.org
We are grateful to Federico Bandi, Luca Benzoni, Jean Jacod, Per Mykland, Roel Oomen, Roberto Renò, Neil Shephard, Viktor Todorov, Lan Zhang, and, in particular Mark Podolskij and Kevin Sheppard for their insights. We also thank participants at the Singapore Management University Conference in Honor of P.C.B. Phillips, July 2008, the CREATES Volatility Symposium, Aarhus, August 2008, the Chicago/London Conference on Financial Markets, December 2008, the Humboldt-Copenhagen Conference on Recent Developments in Financial Econometrics, Berlin, March 2009, the North American Summer Meeting of the Econometric Society, Boston, June 2009, the First European Conference of the Society for Financial Econometrics, Geneva, June 2009, as well as seminar participants at the Federal Reserve Board and the U.S. Commodity Futures Trading Commission for their comments.
Andersen gratefully acknowledges financial support from the NSF through a grant to the NBER and by CREATES funded by the Danish National Research Foundation.
The views in this paper are solely those of the authors and should not be interpreted as reflecting the views of the Board of Governors of the Federal Reserve System, the Federal Reserve Bank of New York or of any other person associated with the Federal Reserve System.

1 Introduction

In the ubiquitous continuous-time no-arbitrage semimartingale framework for modeling asset prices, it is often useful to obtain separate estimates of the continuous part of the volatility process versus the return variation induced by discontinuities or jumps. This is, for instance, the case in the context of risk management, option pricing and volatility forecasting. Thus far, the dominant approach for delivering jump-robust volatility estimates from intradaily return observations has been the so-called realized bipower variation measure, introduced by Barndorff-Nielsen and Shephard (2004), in which volatility is estimated by the cumulative sum of products of adjacent absolute returns.¹ While the bipower variation, by construction, ensures that (finite activity) jumps will not impact the consistency of the volatility estimate, it does not allow for a feasible asymptotic theory under the jump alternative and is subject to a fairly significant finite sample jump distortion (upward bias) that may be of concern in applications. To obtain improved finite sample jump robustness and a feasible asymptotic theory under the jump alternative, the bipower variation has been generalized in subsequent work to tripower and multipower variation measures, which employ products of (lower order) powers of three or more adjacent returns.² The tripower variation is theoretically the most efficient among those. However, it is also more susceptible to market microstructure contamination of the high-frequency return series than the bipower variation. In particular, multipower variation measures are sensitive to the presence of very small (zero) returns arising from stale quotes and rounding to a discrete price grid. In applications, the prevalence of zero returns is often substantial, thus introducing a separate source of potential bias.³

We propose two simple alternatives to the prevailing bipower and tripower variation measures that provide additional robustness to jumps and/or market microstructure noise by using nearest neighbor truncation. The first estimator obtains jump robustness by appropriately scaling the square of the minimum of two consecutive intraday absolute returns. If one of these returns is large, e.g., due to the presence of a jump during the interval, this return is automatically discarded and all weight in the computation falls on the adjacent diffusive returns. Asymptotically, under finite jump activity, we never encounter two adjacent jumps so, like bipower, the measure retains consistency for the underlying integrated diffusive variance. However, this “minimum” or “MinRV” estimator suffers from a similar exposure to small (zero) returns as the traditional multipower variation estimators. In addition, large (absolute) returns are inherently more informative of the underlying volatility

¹Alternative jump-robust estimators include: Mancini (2006), Andersen, Dobrev, and Schaumburg (2008) and Lee and Mykland (2007).

²See Barndorff-Nielsen, Graversen, Jacod, Podolskij, and Shephard (2006) and Barndorff-Nielsen, Shephard, and Winkel (2006).

³Studies of the finite sample behavior of the bipower statistic include Barndorff-Nielsen and Shephard (2004), Huang and Tauchen (2005), Andersen, Bollerslev, and Dobrev (2007), and Dobrev (2007).

than small returns, so our minimum estimator is not particularly efficient. Consequently, we introduce another variant which considers three consecutive intraday returns and simply squares the median absolute return among the three. This estimator also, asymptotically, avoids including the impact of a jump in the measure while reducing the sensitivity to the smallest absolute returns within the trading day and enhancing the asymptotic efficiency.

The unifying theme behind our new estimators is that the absolute returns are truncated at a level controlled by the neighboring returns. “MinRV” uses one-sided truncation as each intraday return is compared only to the subsequent absolute return. The second estimator, denoted “MedRV”, employs two-sided truncation as it uses the median of three adjacent absolute returns. Hence, these estimators exploit an *adaptive* truncation scheme which serves as an endogenous control for the local level of volatility and avoids the potentially delicate choice of an ex-ante threshold required for, e.g., the truncated RV approaches of Mancini (2006) and Aït-Sahalia and Jacod (2007) or the truncated bipower variation of Corsi, Pirino, and Renò (2008).

The endogenous “nearest neighbor” truncation enhances the robustness of our estimators and allows for the development of an asymptotic distribution theory covering both the “no-jump” null hypothesis and the “jump” alternative, which facilitates inference about the presence of jumps. Specifically, the MedRV estimator has better theoretical efficiency properties than the tripower variation measure and displays better finite-sample robustness to jumps and the occurrence of “zero” returns in the sample.

We define the MinRV estimator as arising from the sequential use of the min operator on blocks of two returns and the MedRV estimator from applying the med operator on blocks of three returns. Increasing the block size over this minimum length leads to a gradual efficiency loss, analogous to that observed for higher order multipower variation measures.⁴ Instead, as a theoretically attractive avenue for more efficient jump-robust volatility estimation exploiting larger block sizes, we consider the recent quantile realized volatility (QRV) estimator of Christensen, Oomen, and Podolskij (2008) based on optimally combining relative extreme quantile observations within blocks of twenty or more data points.⁵ However, the reliance on larger blocks has a non-trivial practical cost in finite samples. A critical assumption is that the returns within each block are i.i.d. Gaussian and thus, in particular, that volatility is constant across the block. Although asymptotically valid, this assumption becomes progressively harder to maintain in applied work as the block size increases to encompass a wider calendar interval. This is due to the substantial systematic variation in

⁴Results for such less efficient versions of MinRV and MedRV are available from the authors upon request.

⁵In fact, our “minimum” and “median” estimators share features of both the multipower and quantile estimators. MinRV and MedRV rely on functions of (small) overlapping blocks of adjacent returns like the former, while they exploit the squared quantiles of the (absolute) returns over a (short) block, thus mimicking qualitative aspects of the latter. Another recent estimator of this type is the Realized Outlyingness Weighted Quadratic Covariation (ROWQCov) estimator of Boudt, Croux, and Laurent (2008).

volatility across the trading day, which renders the underlying returns within a longer block non-homogeneous. In addition, although trade and quote arrivals are correlated with increments in volatility they are not well enough aligned to ensure homogeneity (i.e., constant volatility) of the observed log-price increments in either calendar or tick time. Consequently, the gap between the finite sample and asymptotic properties of such estimators tends to be more pronounced than for our “local” MinRV and MedRV estimators. We provide extensive evidence on the finite sample properties of the alternative estimators in simulations and for individual stocks in the Dow-Jones 30 index between January 2005 and May 2008.

The remainder of the paper progresses as follows: Section 2 lays out the basic setup and introduces several popular jump-robust measures of integrated volatility along with our MinRV and MedRV estimators. The asymptotic properties of the new estimators are laid out in a series of propositions. Section 3 provides an empirical application to the set of stocks in the Dow Jones 30 index. Section 4 presents extensive simulation evidence exploring the impact of a variety of features on the performance of the alternative estimators. Section 5 provides concluding observations, while all formal proofs are relegated to the appendix.

2 Jump-Robust IV Estimation

We consider the univariate logarithmic price process $Y = \{Y_t\}_{0 \leq t \leq 1}$ of an asset defined on a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \geq 0}, P)$ so that Y is adapted to the filtration and evolves in continuous time as described by the following jump-diffusive representation,

$$dY_t = \mu_t dt + \sigma_t dB_t + dJ_t \tag{1}$$

where μ is a locally bounded and predictable process and σ is cadlag and bounded away from zero almost surely. The price process is observed at the $N + 1$ discrete points in time $0 \leq t_0 < t_1 < \dots < t_N \leq 1$ over a given period which we refer to as a trading day. The corresponding returns and time intervals are denoted $\Delta Y_i = Y_{t_i} - Y_{t_{i-1}}$ and $\Delta t_i = t_i - t_{i-1}$, $i = 1, \dots, N$. Finally, J denotes a finite activity jump process and dJ_t is either zero (no jump) or a real number indicating the occurrence and size of a jump at time t . Our finite activity jump assumption implies that there are only a finite number of jumps over the trading day. The subsequent analysis is performed via standard continuous record in-fill asymptotics where the time increments between successive return observations, defining the sampling scheme, uniformly shrink towards zero as N increases.

The object of interest is to estimate the continuous part of the quadratic variation, or the integrated variance (IV), defined as

$$IV = \int_0^1 \sigma_u^2 du$$

As is well known, this rules out the use of the popular realized volatility estimator which estimates the total quadratic variation of the observed semimartingale, including the contribution from the cumulative squared jumps.⁶

2.1 Multipower Variation Measures

The initial, and by far most widely used, estimator of IV in the presence of jumps is the bipower variation (BV) measure of Barndorff-Nielsen and Shephard (2004). It can be shown to be consistent for IV in the absence of market microstructure noise but under otherwise very general conditions. It is given as,

$$BV_N = \frac{\pi}{2} \left(\frac{N}{N-1} \right) \sum_{i=1}^{N-1} |\Delta Y_i| |\Delta Y_{i+1}|. \quad (2)$$

The intuition for the consistency and jump robustness of the BV estimator is straightforward: If $\Delta Y_i, \Delta Y_{i+1} \sim i.i.d.N(0, \frac{\sigma^2}{N})$ then $\mathbb{E}[|\Delta Y_i| |\Delta Y_{i+1}|] = \frac{2}{\pi} \frac{\sigma^2}{N}$, and $\left(\frac{N}{N-1} \right)$ is a required finite sample correction factor. As such, each term of the bipower variation measure delivers an unbiased estimate of the underlying local (spot) variance. Moreover, asymptotically, as the returns span near infinitesimal intervals, there will at most be a single jump within two adjacent intervals. This isolated jump will be dampened by the multiplication by a small adjacent (diffusive) return of order $(\frac{1}{\sqrt{N}})$. As N grows this is sufficient to render the jump contribution asymptotically negligible. Nonetheless, in practical applications there will be an upward (finite sample) bias due to large jumps as the adjacent return is not vanishing, reflecting the choice of underlying sampling frequency. The latter is typically governed by market conventions as well as the liquidity and microstructure features of the market. Another drawback of the BV estimator is that the jumps only vanish at the rate of \sqrt{N} which is not sufficient to deliver a continuous-record central limit theory in the presence of jumps in the price path. The desire to obtain an operational asymptotic theory under jump alternatives was a major reason for the introduction of the multipower variation statistics. These are analyzed thoroughly in, e.g., Barndorff-Nielsen, Shephard, and Winkel (2006).

In order to introduce the requisite extension of the bipower variation statistic it is useful, for simplicity, to assume equally spaced sampling, i.e., $\Delta t = t_i - t_{i-1} = 1/N$, for all $i = 1, \dots, N$. The class of multipower variation statistics is then defined via the cumulative

⁶See, e.g., Andersen, Bollerslev, and Diebold (2009), Bandi and Russell (2007), Barndorff-Nielsen and Shephard (2007), McAleer and Medeiros (2008) for surveys of the realized volatility literature.

sum of m products of adjacent absolute returns raised to the (r/m) 'th order, where m is a positive integer and r a positive real number, usually an integer. Hence, the cumulative power of the adjacent products equals r . These statistics provide consistent estimators for the corresponding integrated power of the volatility,

$$MPV_N(m; r) = d_{m,r} \left(\frac{N}{N-m+1} \right) (N)^{r/2-1} \sum_{i=1}^{N-m+1} |\Delta Y_i|^{\frac{r}{m}} \dots |\Delta Y_{i+m-1}|^{\frac{r}{m}} \xrightarrow{P} \int_0^1 \sigma_u^r du \quad (3)$$

where $d_{m,r}$ is a known constant dependent only on m and r , while $\left(\frac{N}{N-m+1} \right)$ is a finite sample correction factor.⁷ If the adjacent returns are i.i.d. Gaussian, each summand in (3) delivers an unbiased estimate of the power of spot volatility. The sum therefore provides a (converging) Riemann approximation to the integrated power of the volatility process.

This multipower variation measure generalizes the entire first generation of estimators in the realized volatility literature, as one obtains the standard realized volatility measure as $RV_N = MPV_N(1; 2)$, while $BV_N = MPV_N(2; 2)$, and additional oft-applied measures include the tripower variation $TPV_N = MPV_N(3; 2)$ the quadpower variation $QPV_N = MPV_N(4; 2)$, and the fourth order power variation $PV_N(4) = MPV_N(1; 4)$. In the presence of a finite activity jump process, the RV estimator is not consistent for the integrated variance, the BV statistic is consistent but does not allow for an asymptotic theory under the jump alternative, while the realized tripower and quadpower measures both provide consistency and allow an associated asymptotic mixed normal limit theory. This property is maintained for $MPV_N(m; 2)$ for $m \geq 3$. Likewise, the fourth order power variation is consistent for the integrated fourth power of the volatility process, the so-called integrated quarticity, but allows for an asymptotic theory only in the absence of jumps. Robust alternatives, which provide both consistency and asymptotic theory under finite activity jumps, are given by $MPV_N(m; 4)$ for $m \geq 5$.⁸

The existence of numerous estimators begs the question of which one is preferable. Not surprisingly, this cannot be answered in general. However, using the ideal setting of no microstructure noise, near infinitely frequent sampling and no jumps, the $MPV_N(m; r)$ measure of lowest order in m delivering the desired feature, whether consistency or a mixed normal limit theory, is the more efficient estimator. Specifically, for estimating the integrated variance in the absence of jumps, the realized volatility estimator is most efficient. Analogously, bipower variation is the preferred consistent jump-robust estimator for IV

⁷In the case of Gaussian i.i.d. price changes $d_{m,r} = \mu_{r/m}^{-m}$, where $\mu_p = E|U|^p = 2^{p/2} \frac{\Gamma(\frac{1}{2}(p+1))}{\Gamma(\frac{1}{2})}$, $U \sim N(0, 1)$, see, e.g., Barndorff-Nielsen and Shephard (2004) and Barndorff-Nielsen, Graversen, Jacod, Podolskij, and Shephard (2006).

⁸See Barndorff-Nielsen, Shephard, and Winkel (2006) for the behavior of the multipower variation estimators under both the no-jump null hypothesis and under the jump alternative in more general scenarios.

while tripower variation is the estimator with minimal asymptotic variance among this class which allows for the development of an asymptotic theory under the jump alternative.

Of course, the frictionless setting is not representative of actual market conditions. In particular, various market features limit the sampling frequency so that the impact of jumps cannot be fully neutralized. In this case, it may be desirable to apply higher order multipower variation measures as they provide better (finite sample) dampening of the jump component. In fact, an extensive simulation exercise by Veraart (2008) finds that the finite sample jump distortion is sufficiently influential to render the $MPV_N(10; 2)$ and $MPV_N(10; 4)$ preferable to the lower order multipower statistics. One caveat is that this simulation evidence assumes a very smooth evolution of the diffusive volatility process and that equally spaced ultra high-frequency returns are available at near arbitrary sampling intervals. These features render volatility near constant across sequences of ten adjacent high-frequency returns thus ensuring that the returns within each block adhere closely to the ideal of being i.i.d. Gaussian distributed. In reality, however, many asset prices display a pronounced U-shape in volatility across the trading day, resulting in sharp movements in volatility across fairly short time intervals. Moreover, fresh quote or trade observations are often not available at the very highest sampling frequencies as both trade and quote intensities undergo significant intraday fluctuations as well. The result is that a block of adjacent (non-stale) price or quote observations often spans a non-trivial time interval, and thus validity of the assumption of homogenous returns and, in particular, constant volatility is questionable. Moreover, the extent of this problem is proportional to the number of adjacent returns exploited by a given estimator. As such, it may be important to explore the impact of the “localness” of the estimator in practical applications. We present evidence both from actual equity data and from an extensive simulation design that this feature, indeed, is a major determinant of the performance of such estimators.

In summary, the class of multipower variation measures embodies a tradeoff between efficiency and localness on the one hand and jump robustness on the other. This motivates our introduction of alternative estimators based on nearest neighbor truncation that retain the local nature of the bipower and tripower estimators and provide better finite sample robustness and asymptotic efficiency, while allowing for feasible asymptotic theory in the presence of (finite activity) jumps.

2.2 The MinRV and MedRV Estimators

We propose the following MinRV and MedRV estimators of integrated variance,

$$\begin{aligned}\text{MinRV}_N &= \frac{\pi}{\pi - 2} \left(\frac{N}{N - 1} \right) \sum_{i=1}^{N-1} \min (|\Delta Y_i|, |\Delta Y_{i+1}|)^2 \\ \text{MedRV}_N &= \frac{\pi}{6 - 4\sqrt{3} + \pi} \left(\frac{N}{N - 2} \right) \sum_{i=2}^{N-1} \text{med} (|\Delta Y_{i-1}|, |\Delta Y_i|, |\Delta Y_{i+1}|)^2\end{aligned}\tag{4}$$

The scaling factors ensure that every summand on the right hand side of equation (4) provides an unbiased estimate of the underlying spot variance if the corresponding block of returns are i.i.d. Gaussian.⁹ The robustness of the MinRV and MedRV estimators compared, e.g., to the BV estimator in equation (2) stems from the fact that returns contaminated by a large jump are fully eliminated by the min/med operators. That is, if a (large) jump occurs within one of the two terms for the MinRV estimator, the min operator simply squares the adjacent (small) diffusive return. There will be an induced bias as we now effectively compute the square of a single return rather than of the minimum of two independent returns, but the bias is an order of magnitude smaller than for bipower. For illustration, assume there is a (large) price jump of size Δj_i in the interval $[t_{i-1}; t_i]$ but no jump in the adjacent intervals so that $|\Delta Y_{i-1}| \ll |\Delta j_i|$ and $|\Delta Y_{i+1}| \ll |\Delta j_i|$. The distortion to the overall BV from the interval containing the jump return clearly dominates the corresponding maximal distortion to the MinRV measure, i.e.,

$$\frac{\pi}{2} |\Delta j_i| (|\Delta Y_{i-1}| + |\Delta Y_{i+1}|) \gg \frac{\pi}{\pi - 2} [|\Delta Y_{i-1}|^2 + |\Delta Y_{i+1}|^2]\tag{5}$$

as the left hand side is of order $(1/\sqrt{N})$ versus order $(1/N)$ on the right hand side. More generally, the upward bias due to jumps for any multipower variation $MPV_N(m; 2)$, $m \geq 2$ is of order $(1/N^{1-\frac{1}{m}})$, thus approaching $(1/N)$ from above for large m . The expression (5) also reflects the important fact that only the number of jumps, but *not* their actual size, have an impact on the bias of the MinRV and MedRV estimators. By construction, one should therefore expect a larger degree of finite sample jump robustness from these estimators than the multipower variations.¹⁰

As indicated above, the MinRV and MedRV estimators are consistent for the integrated variance, as implied by the following proposition.

⁹The derivation of these scaling factors is a simple exercise in integration and is available from the authors upon request.

¹⁰Likewise, the MinRV and MedRV estimators have the advantage of simplicity relative to the truncated RV estimator of Mancini (2006) which truncates jumps above a pre-specified threshold. This threshold choice can be delicate when (latent) volatility is time varying, although some practical progress is reported by Corsi, Pirino, and Renò (2008) using a combination of multipower and threshold techniques.

Proposition 1 *Let the log-price process Y_t be given by the jump-diffusion (1) with finite jump activity. Assume further that μ_t is adapted and locally bounded, σ_t is adapted, cadlag and $\inf_{t \geq 0} \sigma_t > 0$ almost surely. Then we have, as $N \rightarrow \infty$,*

$$\text{MinRV}_N \xrightarrow{P} \int_0^1 \sigma_u^2 du \quad \text{and} \quad \text{MedRV}_N \xrightarrow{P} \int_0^1 \sigma_u^2 du$$

Under slightly stronger assumptions about the volatility process we obtain a corresponding asymptotic distribution theory.

Proposition 2 *Let the log-price process Y_t be given by the jump-diffusion (1) with finite jump activity. Assume further that μ_t is adapted and locally bounded, σ_t is bounded away from zero almost surely and follows an Ito process of the form (A1) given in the appendix, then*

$$\begin{aligned} \sqrt{N}(\text{MinRV}_N - IV) &\xrightarrow{\text{stable } \mathcal{D}} MN \left(0, 3.81 \int_0^1 \sigma_u^4 du \right) \\ \sqrt{N}(\text{MedRV}_N - IV) &\xrightarrow{\text{stable } \mathcal{D}} MN \left(0, 2.96 \int_0^1 \sigma_u^4 du \right) \end{aligned}$$

Moreover, these results remain valid for finite as well as infinite activity jumps in the volatility process subject only to the weak regularity conditions of Barndorff-Nielsen, Graversen, Jacod, Podolskij, and Shephard (2006), stipulating only that the resulting general Ito semimartingale, generalizing Assumption (A1), has jump characteristics that are locally bounded.

The distributional convergence is stable and the notation MN denotes a mixed Gaussian limiting distribution, i.e., a normal distribution conditional on the realization of the integrated quarticity, $IQ = \int_0^1 \sigma_u^4 du$, where, importantly, the limiting normal variate is independent of the (random) IQ process.

The results in Proposition 2 mirror existing limit theories for multipower variation statistics in scenarios without jumps. In this case, the MinRV and MedRV estimators are less efficient than the optimal multipower variation statistics (bipower). Of course, the advantage is the near elimination of finite sample jump distortion.

To illustrate the relationship between the estimators, we derive the joint asymptotic distribution between the MinRV, MedRV and the standard RV and BV estimators under the no-jump null:

Proposition 3 *Under the assumptions of Proposition 2, except that we rule out jumps in the price process, we have the following joint asymptotic stable distributional result,*

$$\sqrt{N} \begin{bmatrix} RV_N - IV \\ BV_N - IV \\ MinRV_N - IV \\ MedRV_N - IV \end{bmatrix} \xrightarrow{\text{stable } \mathcal{D}} MN \left(0, \begin{bmatrix} 2 & 2 & 2 & 2 \\ & 2.61 & 2.98 & 2.53 \\ & & 3.81 & 3.09 \\ & & & 2.96 \end{bmatrix} IQ \right)$$

The proposition shows that there is a non-trivial increase in the asymptotic variance as we move from BV to MinRV, while the MedRV measure is much closer to matching the efficiency of BV. Moreover, both MinRV and MedRV are highly correlated with BV as well as with each other.¹¹ Of course, a more natural comparison is between the MinRV and MedRV estimators and higher order multipower variation measures as these allow for a valid asymptotic theory in the presence of jumps. Table 1 tabulates the relevant efficiency factors. It is evident that MinRV is less efficient, being on par with the sixth order multipower variation estimator. In contrast, MedRV is more efficient (has lower asymptotic variance) than the best (tripower) estimator within this subset of the MPV group.

Proposition 2 implies that feasible inference regarding the realized latent integrated variance is possible based on the MinRV and MedRV estimators, even in the presence of finitely many price jumps. However, as for the MPV measures, this requires a consistent estimator for the integrated quarticity. Such jump-robust estimators are readily constructed from higher order multipower variation statistics. An alternative is to construct an estimator for the quarticity in a direct extension of the principles behind the MinRV and MedRV estimators:

$$\begin{aligned} \text{MinRQ}_N &= \frac{\pi N}{3\pi - 8} \left(\frac{N}{N-1} \right) \sum_{i=1}^{N-1} \min(|\Delta Y_i|, |\Delta Y_{i+1}|)^4 \\ \text{MedRQ}_N &= \frac{3\pi N}{9\pi + 72 - 52\sqrt{3}} \left(\frac{N}{N-2} \right) \sum_{i=2}^{N-1} \text{med}(|\Delta Y_{i-1}|, |\Delta Y_i|, |\Delta Y_{i+1}|)^4 \end{aligned}$$

In fact, the asymptotic theory is entirely analogous and results similar to Propositions 1-3 hold for these quarticity estimators. Likewise, tests for jumps involving statistics based on appropriately normalized differences between the RV and MedRV or MinRV measures may readily be constructed using Proposition 3. We defer the exploration of such procedures to future research.

¹¹We note as a corollary the following correlations between estimators: $\text{Corr}(RV, \text{MinRV}) = 72.4\%$, $\text{Corr}(BV, \text{MinRV}) = 94.5\%$, $\text{Corr}(RV, \text{MedRV}) = 82.2\%$, $\text{Corr}(BV, \text{MedRV}) = 91.0\%$, $\text{Corr}(\text{MinRV}, \text{MedRV}) = 92.0\%$. This suggests that there may be some scope for applying a jump-robust GMM procedure by combining BV, MinRV and MedRV estimators. The derivations are available from the authors upon request.

3 Finite Sample Evidence - Dow Jones 30 Stocks

In this section we gauge the empirical performance of the MinRV and MedRV estimators on the set of Dow Jones 30 stocks using NYSE TAQ data from January 1, 2005 through May 31, 2007. Ignoring short trading days around major holidays we obtain a sample of 601 trading days. We apply the estimators to the series of mid-quotes after filtering out spread outliers (less than 0.1% of the data) and compare the finite sample efficiency of the sub-sampled MinRV and MedRV to sub-sampled multipower variation estimators. Sub-sampling is a simple way of increasing the efficiency of an estimator and was originally advocated for RV by Zhang, Mykland, and Aït-Sahalia (2005). It involves taking the average of an estimator across all possible sub-samples (at a given sampling frequency) obtained by starting from different offsets (and scaling up to match the full day length). For comparison, we also consider a sub-sampled version of the recently developed Quantile RV (QRV) estimator of Christensen, Oomen, and Podolskij (2008). There is little existing evidence regarding the best practical construction of the QRV measure, so we adopt an approach guided by their specific empirical implementation aiming to improve on the efficiency of the bipower variation benchmark.¹²

To keep the exposition manageable we only present results for the various daily measures averaged over the full sample period and across all the stocks. This should help convey the systematic differences across the estimators while reducing the impact of idiosyncratic features of individual stocks and specific time periods.

3.1 “Tick” Time and Calendar Time Sampling

An important aspect of the implementation of any IV estimator is whether to sample in “tick” time or “calendar” time. These sampling schemes represent alternative extreme views of the dependence between observation times and price moves. The tick time approach can be justified (with the standard limiting theory) when volatility is constant in tick time, i.e., observation times and the quadratic variation of the price process are perfectly correlated.

¹²In particular, we use only a single return block for each trading day, consisting of all the relevant recorded quotes, and we optimally weight the quantiles $\{0.05, 0.10, 0.15, 0.85, 0.90, 0.95\}$ based on the finite sample values of the associated scaling factors, exploiting sample sizes all the way up to 23,400 (one per second):

$$QRV = \frac{1}{\nu} \sum_{\lambda \in \{0.85, 0.9, 0.95\}} \omega_{\lambda} [g^2(\lambda) + g^2(1 - \lambda)]$$

where $g(\lambda)$ is the order statistic of the λ^{th} return percentile and ω_{λ} provides the optimal weighting as suggested by Christensen, Oomen, and Podolskij (2008). The constant ν is a normalizing constant whose (finite sample) value must be determined by numerical integration or simulation. Importantly, it is not possible to reliably interpolate the QRV scaling factors across nearby sample sizes due to their pronounced oscillatory nature. This is also the case for a symmetrized version of QRV explored in a conference discussion of Christensen, Oomen, and Podolskij (2008) by Kevin Sheppard.

The calendar time approach, on the other hand, can be justified when observation times are exogenous to the price process, i.e., inference can be carried out conditional on the observation times. In particular, there can be no correlation between quote/trade arrivals and changes in volatility. Neither correlation assumption is likely to be fully accurate, so we study our estimators under both calendar and tick time sampling.¹³

From Figure 1, Panels 1A and 2A we note that the calendar and tick time based approaches lead to qualitatively similar IV estimates for each estimator across the sampling frequencies. A couple of other features are striking. First, it is evident that the RV estimator is more robust than any of the others, as the implied volatility level is quite stable across the entire range of sampling frequencies, spanning 4 seconds to 5 minutes. In contrast, all the jump-robust IV estimators vary significantly across the frequencies, with the pronounced dip at the highest frequencies being particularly noteworthy.¹⁴ In most cases, the maximum value is obtained around the two-minute frequency. The subsequent explorations document fairly significant downward biases at the highest and lowest frequencies, suggesting that IV estimates based on sampling around the two-minute level are the least biased. In particular, we study the effects of stale quotes, unevenly spaced data, i.i.d. microstructure noise and strong intraday volatility patterns below. Second, the IV estimators differ substantially from each other: MinRV, MedRV and BV lead to roughly similar average estimates while the tripower (TV) and the quantile (QRV) estimators often yield much lower and RV much higher estimates. These differences are highly significant given the standard deviation of daily IV estimates depicted in Figure 2, Panels 1A-2A, taking into account the large number of underlying days and stocks. For RV, the higher estimates are consistent with the presence of price jumps as, by construction, it is not jump-robust. As such, the discrepancy between RV and the other measures provides an indication of the overall jump contribution to the return variation across the Dow Jones stocks. The analysis in the following sections sheds further light on the reasons for the disparate behavior of the alternative IV measures.

3.2 Stale Quotes

Table 2 summarizes basic descriptive statistics for the Dow Jones components, including all companies that were part of the index during January 1, 2005 and May 31, 2007. These stocks are generally very actively traded, with an average of one new quote arrival every 2 seconds throughout the normal trading hours from 9:30am to 4:00pm EST. However, a

¹³In practice, this issue seems greatly alleviated if one avoids sampling at ultra-high frequencies. For example, the effect appears empirically negligible for actively traded securities if volatility is computed from one-minute returns. Nonetheless, much current research aims to reduce the impact of noise sufficiently that even the highest sampling frequencies may be exploited productively. As such, the dependence between observation times (quotes or trades) and the underlying price process is an intriguing area for future research. See Mykland, Renault, and Zhang (2008) for a recent discussion of volatility estimation in such circumstances.

¹⁴Chaboud, Chiquoine, Hjalmarsson, and Loretan (2007) have also documented a similar dip of the bipower variation estimates of IV on FX and T-Bond data.

significant number of these quotes simply repeat previous quotes, as reported in the first two columns of the table: the average number of non-duplicate quotes is only about one quarter of the recorded quotes. Remarkably, the two most actively traded stocks (INTC and MSFT) have the smallest number of non-duplicate quotes. As a consequence, estimators may well behave differently if based on duplicate-filtered versus unfiltered data. Furthermore, the last column of the table indicates that the duplicate quotes are clustered as most stocks experience long spells of duplicate quotes. The median number of zero returns lasting longer than 30 seconds on each day is 161, those lasting longer than 1 minute are about 48 on each day, and those lasting longer than 2 minutes are about 11 per day. Importantly, this pattern is observed without much variation across stocks (reported) or trading days (not reported). The effect of filtering out duplicates is shown in Figure 1, Panel 3A. For all estimators, except QRV, the dip at higher frequencies is substantially alleviated without any major impact at lower frequencies, consistent with the elimination of a bias induced by zero returns. Specifically, we note that the behavior of the TV measure now starts to resemble that of the remaining IV estimators, leaving QRV as a relative outlier.

3.3 Microstructure Noise

The prevalent way of dealing with microstructure noise is to sample at a lower frequency than the available data (e.g., 2 minutes) to gain noise robustness and then compensate for the efficiency loss by subsampling the estimator. An alternative approach termed “pre-averaging,” recently introduced by Podolskij and Vetter (2006), exploits the data at the highest frequency available, but uses local “pre-averaging” via a kernel function to produce a set of non-overlapping (asymptotically) noise free (in practice, noise reduced) observations to which standard IV estimators may be applied. In practice, this necessitates a choice of bandwidth (and kernel function) and leads to a familiar bias-variance trade-off which we study for our set of estimators. To render the results comparable, we consider pre-averaging window lengths matching our sub-sampling frequencies. In particular, we note that if the pre-averaged log-prices for window of length K are defined as $\bar{Y}_i = \frac{1}{K} \sum_{j=0}^{K-1} Y_{i+j}$, $i = 0, 1, \dots, N - K + 1$ then the pre-averaged version of each estimator has equivalent representation in terms of sub-sampling \bar{Y}_i at frequency K observations, implying that the corresponding returns $\bar{Y}_{i+K} - \bar{Y}_i$ cover a window of length $2K$ observations of the original log-price series Y_i . Therefore, for each pre-averaged IV estimator based on \bar{Y}_i we set the underlying pre-averaging window length K , so that $2K$ equals the sample frequency of each corresponding sub-sampled IV estimator based on Y_i .¹⁵

¹⁵Unlike the sub-sampled IV estimators, their pre-averaged counterparts require an additional scaling factor implied by the kernel function and asymptotically equal to 3, see, e.g., Podolskij and Vetter (2006) and Christensen, Oomen, and Podolskij (2008). We computed the exact finite sample value of this scaling factor (available upon request) as a function of pre-averaging window length in order to eliminate the bias otherwise incurred when using its asymptotic value.

The pre-averaged estimators are displayed in Figure 1, Panels 1B-3B. Overall, they mirror the qualitative behavior of the sub-sampled estimators although the estimated level of volatility is marginally lower. In particular, the tripower variation measure continues to be strongly downward biased. Moreover, the QRV measure again delivers values that fall at the bottom of the range across the estimators. This goes along with a substantially lower daily standard deviation of the measure as depicted in Figure 2. The simulation section below exemplifies some of the features that may rationalize this distinctive behavior of QRV.

3.4 Robustness across Volatility Regimes

Based on newly available NYSE/TAQ data, we consider a second shorter sample covering June 1, 2007 to May 31, 2008. Ignoring short trading days around major holidays we obtain a sample of 248 days during the second sample period. This allows us to consider the robustness and relative performance of the estimators across different regimes as volatility is significantly higher during the more recent period.

The results are displayed in Figures 3-4. Along with the much higher estimates of IV there is also a much wider gap between RV and the jump-robust measures suggesting more significant jump activity in this period. The relative behavior of the individual estimators, however, follows roughly the same pattern as in the first sample, with notably lower estimates of QRV and TV compared to MinRV, MedRV and BV. Interestingly, Panels 3A and 3B also suggest that, in this period of higher volatility, it may be particularly beneficial to filter out duplicates for the sake of producing volatility estimates that are consistent across a wider range of sampling frequencies and pre-averaging windows.

Overall, the behavior of the various IV estimators appears robust to the volatility regime as the qualitative differences observed over the prior sample remain intact. It is worth noting, though, that the dispersion of the estimates is no longer wildly inflated at the highest frequencies as was the case in the initial sample period (Figure 4 vs. Figure 2). This may reflect a reduction in the noise-to-signal ratio over time due to the growing quantity and quality of the high-frequency quotations.¹⁶

4 Finite Sample Simulation Evidence

We conduct Monte Carlo experiments focusing on features of the data generating process that may affect the finite sample behavior of the various IV estimators. In particular, we compare the performance of sub-sampled/pre-averaged MinRV and MedRV estimators to sub-sampled/pre-averaged BV, TV, and QRV benchmarks for a set of models embodying distinct features. The emphasis is on the qualitative impact. In reality, all these features are

¹⁶In addition, a period of elevated volatility, without a commensurate increase in the microstructure noise, translates into an improved signal-to-noise ratio, i.e. better relative quality of the data.

likely present simultaneously and interact with each other, creating rather complex patterns in tick-by-tick data. Hence, the simulations are not designed to replicate the quantitative magnitude of statistics observed from actual data in all dimensions, but rather to help identify features which may be relevant to explain the systematic patterns in the empirical results.

We consider the following set of volatility models, which mostly deviate from the simple i.i.d. Gaussian benchmark in a single dimension to facilitate direct interpretation of the qualitative impact:

- *Model 1*: “BM”. This is our baseline Brownian motion model with sampling on an equispaced time grid. It provides the ideal setting under which the finite sample performance of the IV estimators should be closely in line with the underlying asymptotic theory.
- *Model 2*: “SV-U”. This is a stochastic volatility model (two-factor affine) with intraday U-shape volatility pattern and sampling on an equispaced time grid. It allows us to isolate potential finite sample biases of the estimators due to time variation in volatility.
- *Model 3*: “BM + Sparsity”. This is a Brownian motion model with sampling on a sparse (exogenously random) time grid. While not necessarily realistic, this model is helpful for studying the potential distortion of the estimators when applied on non-homogeneously sampled returns, effectively inducing spurious variations in their volatility.
- *Model 4*: “BM + 1 Jump”. This is a Brownian motion model with one jump on each day and sampling on an equispaced time grid. It serves to illustrate the degree of finite sample jump robustness of the alternative IV estimators.
- *Model 5*: “BM + 4 Jumps”. This is a Brownian motion model with four jumps on each day and sampling on an equispaced time grid. We use this jump specification to study the impact of multiple (potentially adjacent) jumps.
- *Model 6*: “BM + Noise”. Brownian motion model with sampling on an equispaced time grid and subject to i.i.d. microstructure noise. It allows us to shed light on potential distortions due to microstructure noise.

4.1 Simulation Design

In each model, the price process $\{Y_t\}$ follows a driftless Brownian motion with instantaneous volatility $\sigma(t)$:

$$dY(t) = \sigma(t) dW_1(t)$$

Across all model specifications, the unconditional IV of each day is calibrated to 0.000159 corresponding to an annualized volatility of 20% (assuming 252 trading days per year). This roughly matches the average level of volatility observed in our DJ 30 sample between January 2005 and May 2007. We also match the average sample frequency of 2 seconds

resulting in 11,700 intraday observations. We adopt an equispaced time grid across all models except for Model 3, where we use (exogenously) random sampling.

In Model 2, the stochastic volatility model with intraday U-shape volatility pattern is described by

$$\begin{aligned} dY(t) &= \sigma_u(t) \sigma_{sv}(t) dW_1(t) \\ \sigma_{sv}^2(t) &= \sigma_1^2(t) + \sigma_2^2(t) \\ d\sigma_1^2(t) &= \kappa_1 [\theta_1 - \sigma_1^2(t)] dt + \eta_1 \sigma_1(t) dW_{21}(t) \\ d\sigma_2^2(t) &= \kappa_2 [\theta_2 - \sigma_2^2(t)] dt + \eta_2 \sigma_2(t) dW_{22}(t) \end{aligned}$$

with W_{21}, W_{22} independent and the leverage effect captured by the instantaneous correlations $\rho_1 = \text{corr}(dW_1(t), dW_{21}(t))$ and $\rho_2 = \text{corr}(dW_1(t), dW_{22}(t))$. The two factor model parameters are calibrated in line with Andersen, Bollerslev, and Meddahi (2005) in percentage form as $\kappa_1 = 0.6$, $\kappa_2 = 0.1$, $\theta_1 = 1.0582$, $\theta_2 = 0.5291$, $\eta_1 = 0.2$, $\eta_2 = 0.1$, $\rho_1 = 0.9$, $\rho_2 = -0.4$. Following Hasbrouck (1999) we model the diurnal volatility U-shape as the sum of two exponentials:

$$\sigma_u(t) = C + A e^{-at} + B e^{-b(1-t)}, \quad t \in [0; 1]$$

where the constants $A = 0.75$, $B = 0.25$, $C = 0.88929198$, $a = 10$, $b = 10$ are calibrated to produce a strong asymmetric U-shape with variance at the open ($t = 0$) more than 3 times the midday variance ($t = 1/2$) and variance at the close about 1.5 times the midday variance.

In Model 3, non-homogeneous sampling is obtained by taking a random sample of 11,701 points (without repetition) out of the full daily time grid (23,401 seconds from 9:30 am to 4:00 pm). The resulting sample size is identical to the one obtained with 2-second equidistant sampling, but the price observations are subject to exogenously imposed random sparsity.

In Models 4-5, price jumps are introduced by extending the log-price process as

$$dY(t) = \sigma(t) dW_1(t) + dJ_t$$

where the Poisson jump process (J_t) is assumed independent of (W_1, W_2) . In order to stress test the IV estimators, we calibrate the process to match moderate “jump days” on which one (Model 4) or four (Model 5) Gaussian jumps account for an average increase of 25% in realized volatility (i.e. the jump contribution JV is 25% of IV or, equivalently, 20% of QV = IV + JV, thereby generating many jumps that are not too obvious return outliers).

Finally in Model 6, we simulate Gaussian i.i.d. noise with a moderate noise-to-signal ratio $\lambda = 0.25$, defined as the ratio of annualized error variance to annualized IV.

4.2 Simulation Results

For each model specification, we simulate 2,500 trading days (roughly ten years of data ignoring non-trading periods) and tabulate the relative bias and efficiency of the sub-sampled IV estimators at sampling frequencies 12, 60, and 300 seconds (Table 3) as well as their pre-averaged counterparts for pre-averaging windows 12, 60, and 300 seconds (Table 4).¹⁷

The results for Model 1, Tables 3-4, confirm that, in a frictionless setting with homogeneous returns, all IV estimators are unbiased and their relative efficiency is closely in line with the asymptotic theory. However, under the more realistic scenario of stochastic volatility and pronounced diurnal volatility patterns all IV estimators exhibit a pronounced downward bias if sub-sampled or pre-averaged sparsely (bottom panels, Tables 3-4, Model 2). Moreover, QRV remains biased even for relatively high sub-sampling frequencies or small pre-averaging windows (top panels, Tables 3-4, Model 2), whereas the remaining IV estimators become unbiased as the sampling frequency grows as they are better equipped to handle intraday volatility fluctuations due to the “locality” achieved by using short rolling blocks of returns. Thus, the MinRV/MedRV and the multipower variation measures seem to have a clear finite sample advantage over QRV in this regard.

At the same time, all estimators are downward biased at the highest frequencies if the sampled returns are non-homogeneous due to (exogenous) random sparsity of the available observations (top panels, Tables 3-4, Model 3). In this case, pre-averaging over wider windows or sampling at sparser frequencies (bottom panels, Tables 3-4, Model 3) eliminates the bias for all estimators. On the other hand, the results for the jump scenarios (Models 4-5, Tables 3-4) provide evidence that MinRV, MedRV, and QRV are considerably less biased in the presence of jumps, especially multiple ones, compared to the multipower variation measures. Finally, the sensitivity of the estimators to microstructure noise is quite similar and both sub-sampling (middle and bottom panels, Table 3, Model 6) and pre-averaging (middle and bottom panels, Table 4, Model 6) seem to offer a sensible solution for all estimators. It is worth noting, though, that matching the pre-averaging window size to the sub-sampling frequency results in consistently lower MSE of the pre-averaged estimators compared to their sub-sampled counterparts (Table 3 vs Table 4), which is in line with the underlying asymptotic theory.

These controlled Monte Carlo experiments suggest that the MinRV/MedRV estimators combine the main advantages of the existing IV estimators in finite samples: superior robustness to jumps akin to QRV as well as reasonable robustness to time-varying volatility

¹⁷The relative bias is computed as the sample mean of \widehat{IV}/IV , while the relative efficiency factor at the 60-sec sample frequency is calculated as the sample mean of $390(\widehat{IV} - IV)^2/IQ$, where IV and IQ are the true simulated integrated variance and integrated quarticity on each day, while \widehat{IV} is the IV estimate for the given day. For example, the MSE factor for 60-sec sub-sampled RV is about 1.33, and thus in line with the theoretical value derived in Zhang, Mykland, and Aït-Sahalia (2005).

like the multipower variation measures. Sub-sampling or pre-averaging either too sparsely (i.e. much lower than one-minute frequency) or too frequently (much higher than one-minute frequency) result in downward biases for all the IV estimators, but especially in the case of QRV, if there are pronounced intraday variations in volatility or instances of spurious sparsity of the price observations. In this regard, the evidence in Tables 3-4 is qualitatively consistent with the signature plots produced for the Dow Jones 30 stocks in Section 3 (Figures 1 and 3). As such, the return variation estimates obtained around the 2-minute frequency appear most reliable as they seem to avoid both the (downward) biases induced at the highest frequencies by sparsity, noise and rounding and those at lower frequencies due to stochastic volatility and diurnal patterns.

5 Conclusion

We introduce two new jump-robust estimators of integrated variance based on high-frequency return observations. These MinRV and MedRV estimators rely on nearest neighbor truncation as an attractive novel way to achieve jump robustness, while sharing a number of important features with existing estimators. First, the estimators mirror the traditional RV measure in simply cumulating a sum of squared intraday returns. Second, they resemble the bipower and tripower variation estimators in exploiting two or three adjacent return observations to obtain each summand within the sum defining the variation measure. An important distinction, however, is that the MinRV and MedRV estimators dampen the effect of jumps at a faster asymptotic rate than any multipower variation estimator. Third, they provide an even sharper truncation of outliers than the threshold RV estimators. The main difference is that MinRV and MedRV exploit a threshold given by the adjacent return observations which ensures a local and adaptive truncation level. In contrast, threshold RV estimators determine the appropriate threshold by some auxiliary procedure and require effort to ensure sensible adaptation to the time-varying return volatility level. Fourth, since the minimum and the median correspond to particular quantiles, the MinRV and MedRV estimators are conceptually related to quantile RV estimators although there are several important differences. The quantile RV estimators exploit non-overlapping blocks and emphasize the efficiency gains from using extreme quantiles based on long blocks of observations, thereby sacrificing the localness of the estimator. The MinRV and MedRV achieve good efficiency properties exploiting very short overlapping blocks. Moreover, the finite sample scaling factors and normalizing constants of the MinRV and MedRV are known in closed form and do not require costly numerical evaluation.

The MinRV and MedRV estimators are designed to obtain a number of specific properties. One, the finite sample jump-robustness is, by construction, excellent as influential (large) jumps are systematically discarded. Two, the use of overlapping windows helps

extract additional information from the observed data. Three, the local nature of the estimators renders them robust to strong intraday variation in volatility. Four, the adaptive nature of the (implicit) threshold avoids any delicate calibration of the appropriate cut-off level. Five, there is no need for auxiliary procedures to determine a window length or bandwidth. Six, the MedRV estimator is robust also to the presence of spurious zero returns (quote or trade price duplicates). Finally, both MinRV and MedRV are very simple and entail only minor modifications of the popular realized volatility, bipower and tripower variation measures. In particular, the proofs of the asymptotic theory for the estimators, although rather technical, can be derived in a manner quite similar to those in the extant literature. In fact, our appendix illustrates how one may recast the features of a specific estimator into the format required for application of the existing powerful methodology for deriving the asymptotic distribution theory.

The evidence gleaned from our analysis of the Dow Jones 30 stocks as well as the simulation study confirms that the MinRV and MedRV estimators possess excellent jump robustness (only minor upward bias), while with an adequate choice of a sampling frequency they also may be designed to avoid significant downward biases from sparse data and/or zero returns. Moreover, the MedRV measure is theoretically more efficient than all existing jump-robust multipower variation estimators which allow for development of an asymptotic limit theory in the presence of jumps, i.e., the tripower and higher order power variation measures. In practice, MinRV and MedRV appear to perform on par with the subsampled bipower variation statistic, while clearly improving on the jump-robustness of the latter. Finally, the new estimators appear to dominate the less local quantile RV estimators as the latter tend to suffer significantly from the presence of a strong and systematic intraday variation in the volatility process and other factors inducing violations of return homogeneity across larger blocks of data.

In conclusion, the MinRV and MedRV measures are promising candidates for practical applications involving the estimation of integrated variance due to their combination of reasonable efficiency and good robustness properties. Such estimators may be particularly attractive for estimation and inference in settings where the presence of jumps cannot be ignored. Moreover, it is simple to generalize these estimators to obtain corresponding measures of the integrated quarticity and the associated asymptotic limit theory is straightforward to derive using the proof strategy developed in this paper. As an example, the estimators should be useful ingredients in procedures designed to test for the presence of price jumps.

References

- Aït-Sahalia, Y. and J. Jacod (2007). Volatility estimators for discretely sampled lévy processes. *Annals of Statistics* 35(1), 355–392.
- Andersen, T. G., T. Bollerslev, and F. X. Diebold (2009). *Parametric and nonparametric volatility measurement*. North Holland. In Handbook of Financial Econometrics, Yacine Aït-Sahalia, Lars P. Hansen, and Jose A. Scheinkman (Eds.).
- Andersen, T. G., T. Bollerslev, and D. Dobrev (2007). No-arbitrage semi-martingale restrictions for continuous-time volatility models subject to leverage effects, jumps and i.i.d. noise: Theory and testable distributional implications. *Journal of Econometrics* 138(1), 125–80.
- Andersen, T. G., T. Bollerslev, and N. Meddahi (2005). Correcting the errors: Volatility forecast evaluation using high-frequency data and realized volatilities. *Econometrica* 73(1), 279–96.
- Andersen, T. G., D. P. Dobrev, and E. Schaumburg (2008). Duration based volatility estimation. *Manuscript, Northwestern University*.
- Bandi, F. M. and J. R. Russell (2007). *Volatility*. Elsevier Science, New York. in Handbook of Financial Engineering, V. Linetski and J. Birge (Eds.).
- Barndorff-Nielsen, O. E., S. E. Graversen, J. Jacod, M. Podolskij, and N. Shephard (2006). A central limit theorem for realized power and bipower variations of continuous semimartingales. in *From stochastic calculus to mathematical finance : the Shiryaev Festschrift*, Springer Verlag.
- Barndorff-Nielsen, O. E., S. E. Graversen, J. Jacod, and N. Shephard (2006). Limit theorems for bipower variation in financial econometrics. *Econometric Theory* 22(4), 677–719.
- Barndorff-Nielsen, O. E. and N. Shephard (2004). Power and bipower variation with stochastic volatility and jumps. *Journal of Financial Econometrics* 2(1), 1–37.
- Barndorff-Nielsen, O. E. and N. Shephard (2007). *Variation, Jumps, Market Frictions and High Frequency Data in Financial Econometrics*. Cambridge University Press. In Advances in Economics and Econometrics. Theory and Applications, Ninth World Congress, R. Blundell, T. Persson, and W. Newey (Eds.).
- Barndorff-Nielsen, O. E., N. Shephard, and M. Winkel (2006). Limit theorems for multipower variation in the presence of jumps. *Stochastic Processes and Their Applications* 116, 796–806.
- Boudt, K., C. Croux, and S. Laurent (2008). Outlyingness weighted quadratic covariation. *Working paper*.
- Chaboud, A., B. Chiquoine, E. Hjalmarsson, and M. Loretan (2007). Frequency of observation and the estimation of integrated volatility in deep and liquid markets. *Board of Governors of the Federal Reserve System, International Finance Discussion Papers*.
- Christensen, K., R. Oomen, and M. Podolskij (2008). Realized quantile-based estimation of integrated variance. *Working paper*.
- Corsi, F., D. Pirino, and R. Renò (2008). Volatility forecasting: The jumps do matter. *SSRN Working Paper*.
- Dobrev, D. (2007). Capturing volatility from large price moves: generalized range theory and applications. *Working paper, Northwestern University*.

- Hasbrouck, J. (1999). The dynamics of discrete bid and ask quotes. *Journal of Finance* 54(6), 2109–42.
- Huang, X. and G. Tauchen (2005). The relative contribution of jumps to total price variance. *Journal of Financial Econometrics* 3(4), 456–99.
- Jacod, J. and A. N. Shiryaev (2003). *Limit theorems for stochastic processes* (2nd ed.). Grundlehren der mathematischen Wissenschaften. Berlin; New York: Springer.
- Lee, S. S. and P. A. Mykland (2007, December). Jumps in financial markets: A new non-parametric test and jump dynamics. *Review of Financial Studies* 20.
- Mancini, C. (2006). Estimating the integrated volatility in stochastic volatility models with levy type jumps. *Working paper, University of Firenze*.
- McAleer, M. and M. C. Medeiros (2008). Realized volatility: A review. *Econometric Reviews* 27(1-3), 10–45.
- Mykland, P. A., E. Renault, and L. Zhang (2008). Volatility estimation when trade times and returns are dependent. *Working paper University of Chicago and UIC*.
- Mykland, P. A. and L. Zhang (2009). Inference for continuous semimartingales observed at high frequency. *Econometrica* 77(5), 1403–1445.
- Podolskij, M. and M. Vetter (2006). Estimation of volatility functionals in the simultaneous presence of microstructure noise and jumps. *Working paper, CREATES, University of Aarhus and Ruhr University of Bochum*.
- Revuz, D. and M. Yor (1999). *Continuous martingales and Brownian motion* (3rd ed.). Grundlehren der mathematischen Wissenschaften. Berlin [u.a.]: Springer.
- Veraart, A. E. D. (2008). Inference for the jump part of quadratic variation of itô semimartingales. *working paper University of Aarhus*.
- Zhang, L., P. A. Mykland, and Y. Aït-Sahalia (2005). A tale of two time scales: Determining integrated volatility with noisy high-frequency data. *Journal of the American Statistical Association* 100(472), 1394–1411.

A Consistency and CLT

Let Y_t be the log price process and assume that it follows a Brownian semimartingale

$$Y_t = Y_0 + \int_0^t a_u du + \int_0^t \sigma_{u-} dB_u \quad (6)$$

where a is a locally bounded and predictable process and σ is adapted and cadlag and bounded away from zero. Without loss of generality, we further assume that the functions a, σ are uniformly bounded and that $\inf_{t>0} \sigma_t > 0$ a.s.¹⁸ The extension allowing for finite activity jumps in Y_t is dealt with in Section A.3 below.

For the central limit theorem we require in addition that the volatility process follows a generalized Itô process:

$$\textbf{Assumption (A1)} : \sigma_t = \sigma_0 + \int_0^t \tilde{a}_u du + \int_0^t \tilde{\sigma}_{u-} dB_u + \int_0^t \tilde{v}_{u-} dW_u,$$

where \tilde{a} is locally bounded and predictable and $\tilde{\sigma}, \tilde{v}$ are cadlag and the Brownian motions B, W are uncorrelated. As before, we impose without loss of generality that the functions $\tilde{a}, \tilde{\sigma}$, and \tilde{v} are uniformly bounded and that $\inf_{t>0} \tilde{\sigma}_t > 0$ and $\inf_{t>0} \tilde{v}_t > 0$ a.s. In addition, as explained at the end of Section A.2, a general set of jump processes may be included in the volatility process specification without altering the results.

We assume that Y is observed at $N + 1$ evenly spaced time points spanning the interval $[0; 1]$. Below, we denote these observations by $Y_{i/N}$, $i = 0, \dots, N$, and the associated log-returns by $\Delta_i^N Y = Y_{i/N} - Y_{(i-1)/N}$, $i = 1, \dots, N$. The proofs involve sequences of standardized return observations and corresponding approximating sequences for which volatility is fixed across one or more returns. Hence, we introduce non-overlapping blocks of $M \geq 1$ returns for which the volatility process is constant. We assume we have $K = N/M$ such blocks in the sample. Consequently, we define the quantities,

$$\chi_i^N = \sqrt{N} \Delta_i^N Y \quad \text{and} \quad (7)$$

$$\beta_i^{N,M} = \sqrt{N} \sigma_{\lfloor (i-1)/M \rfloor / N} \Delta_i^N Y = \sqrt{N} \sigma_{\lfloor (i-1)/M \rfloor / K} \Delta_i^N Y, \quad (8)$$

where $\lfloor \cdot \rfloor$ indicates the integer part of an expression. Hence, for each of the K return blocks, corresponding to $\beta_i^{N,M}$, the volatility remains fixed at the value it attains at the beginning of the block. We shall exploit that, for large N , $\chi_i^N \approx \beta_i^{N,M}$. The strategy of the proof is then, as in Barndorff-Nielsen, Graversen, Jacod, Podolskij, and Shephard (2006), henceforth BNGJPS, to first show convergence in probability and distribution for the approximate process and then argue that the difference is small.

Let $g : \mathbb{R}^2 \mapsto \mathbb{R}_+$ be given by,

$$g(x) = \frac{\pi}{\pi - 2} \min(|x_1|^2, |x_2|^2).$$

then, for any two bivariate vectors, $\mathbf{a} = (a_1, a_2)$ and $\mathbf{b} = (b_1, b_2)$, we have the bound

$$|g(\mathbf{a}) - g(\mathbf{b})| \leq (|a_1^2 - b_1^2| + |a_2^2 - b_2^2|) \quad (9)$$

¹⁸As argued in Barndorff-Nielsen, Graversen, Jacod, Podolskij, and Shephard (2006), this follows from working with the stopped versions of the processes: $T_t^{(k)} = Y_{t \wedge T_k}$ and $\sigma_t^{(k)} = \sigma_{t \wedge T_k}$ where $T_k = \inf\{t | |a_t| + |\sigma_{t-}| \geq k\}$ and $T_k \nearrow \infty$ a.s.

and furthermore we note that $g(x, y)$ is differentiable, except on the null set $\{(x, y) \in \mathbb{R}^2 | x = y\}$ and

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} [\min(x_1^2, x_2^2 + \varepsilon z) - \min(x_1^2, x_2^2)] = \begin{cases} z & \text{if } x_2^2 < x_1^2 \\ 0 & \text{if } x_2^2 > x_1^2 \end{cases} \quad (10)$$

Our MinRV estimator of IV takes the form,

$$\text{minRV}_N = \frac{\pi}{\pi - 2} \frac{1}{N-1} \sum_{i=1}^{N-1} \min((\chi_i^N)^2, (\chi_{i+1}^N)^2) = \frac{1}{N-1} \sum_{i=1}^{N-1} g(\chi_i^N, \chi_{i+1}^N).$$

The proof revolves around the sequences,

$$\mathbf{V}_N = \frac{1}{N} \sum_{i=1}^{N-1} g(\chi_i^N, \chi_{i+1}^N) \quad \text{and} \quad \mathbf{U}_N^M = \frac{1}{N} \sum_{i=1}^{N-1} g(\beta_i^{N,M}, \beta_{i+1}^{N,M}).$$

Since $\text{MinRV}_N = \frac{N}{N-1} \mathbf{V}_N$ the \mathbf{V}_N sequence is asymptotically equivalent to our MinRV estimator, while \mathbf{U}_N^M is the approximating sequence.

We introduce some additional simplifying notation: for any adapted, integrable, d -dimensional cadlag process, Z , and for $N \geq j > i-1 \geq 0$ we define the expectation conditional on information at time $\frac{i-1}{N}$:

$$E_{i-1} \left[Z_{\frac{j}{N}} \right] = E \left[Z_{\frac{j}{N}} | \mathcal{F}_{\frac{i-1}{N}} \right] \quad (11)$$

One useful implication of our ability to focus on the case with uniformly bounded drift and volatility functions is that, using the Burkholder-Davis-Grundy inequalities, we have,

$$E_{i-1} [|\chi_i^N|^p] \leq C \quad \text{and} \quad E_{i-1} [|\beta_i^{N,M}|^p] \leq C, \quad (12)$$

where $p > 0$ and C denotes a generic positive constant which (with a slight abuse of notation) will take on disparate values in different places in what follows. We exploit this property of uniformly bounded moments repeatedly in the sequel.

We may now decompose our basic estimators for IV into a sum of conditional expectations and the associated martingale difference sequence: $\mathbf{V}_N = \mathbf{V}_{1N} + \mathbf{V}_{2N}$ and $\mathbf{U}_N^M = \mathbf{U}_{1N}^M + \mathbf{U}_{2N}^M$ where,

$$\begin{aligned} \mathbf{V}_{1N} &= \frac{1}{N} \sum_{i=1}^{N-1} E_{i-1} [g(\chi_i^N, \chi_{i+1}^N)], & \mathbf{V}_{2N} &= \frac{1}{N} \sum_{i=1}^{N-1} \{g(\chi_i^N, \chi_{i+1}^N) - E_{i-1} [g(\chi_i^N, \chi_{i+1}^N)]\} \\ \mathbf{U}_{1N}^M &= \frac{1}{N} \sum_{i=1}^{N-1} E_{\lfloor \frac{i-1}{M} \rfloor M} [g(\beta_i^{N,M}, \beta_{i+1}^{N,M})], & \mathbf{U}_{2N}^M &= \frac{1}{N} \sum_{i=1}^{N-1} \{g(\beta_i^{N,M}, \beta_{i+1}^{N,M}) - E_{\lfloor \frac{i-1}{M} \rfloor M} [g(\beta_i^{N,M}, \beta_{i+1}^{N,M})]\}. \end{aligned}$$

When $M = 1$ we will use the shorthand $\beta_i^N \equiv \beta_i^{N,1}$, $\mathbf{U}_N \equiv \mathbf{U}_N^1$ and similarly for the individual pieces \mathbf{U}_{1N} and \mathbf{U}_{2N} . These definitions allow us to decompose the main estimator:

$$\mathbf{V}_N = \mathbf{U}_{1N} + \mathbf{U}_{2N} + (\mathbf{V}_{1N} - \mathbf{U}_{1N}) + (\mathbf{V}_{2N} - \mathbf{U}_{2N}) \quad (13)$$

Consistency of \mathbf{V}_N can then be obtained by showing consistency of the estimator applied to the approximating Brownian path with piecewise constant volatility ($\mathbf{U}_N = \mathbf{U}_{1N} + \mathbf{U}_{2N}$) and then showing that the difference $\mathbf{V}_N - \mathbf{U}_N$ (the last two terms) is asymptotically negligible. This is what we do in Section A.1 below. To

prove a CLT, we exploit a different decomposition (similar to Mykland and Zhang (2009)), in which we show the CLT for our estimator applied to an approximating Brownian motion in which volatility is held constant over a block of length M and then proceed to show that the difference between the original estimator and the estimator applied to the approximating process is negligible. This analysis is carried out in Section A.2 based on the decomposition:

$$\begin{aligned}\sqrt{N}(\mathbf{V}_N - IV) &= \sqrt{N}(\mathbf{V}_{1N} - \mathbf{U}_{1N}) + \sqrt{N}(\mathbf{V}_{2N} - \mathbf{U}_{2N}) + \sqrt{N}(\mathbf{U}_{1N} - IV) \\ &+ \sqrt{N}(\mathbf{U}_{2N} - \mathbf{U}_{2N}^M) + \sqrt{N}\mathbf{U}_{2N}^M\end{aligned}\quad (14)$$

A.1 Proposition 1: Consistency

We proceed by analyzing equation (13) term by term through a series of lemmas. For brevity, we focus on the features that are specific to our estimator, while referring to proofs in the extant literature when feasible. This also serves to highlight the underlying structural similarities between our IV measure and some previously proposed IV estimators.

Lemma 4 *Under the maintained assumptions we have,*

$$\mathbf{U}_{1N} \xrightarrow{P} IV \quad (15)$$

Moreover, if Assumption (A1) holds we obtain,

$$\sqrt{N}(\mathbf{U}_{1N} - IV) \xrightarrow{P} 0 \quad (16)$$

Proof. First, note that

$$g(\beta_i^N, \beta_{i+1}^N) = \left[g(\beta_i^N, \beta_{i+1}^N) - g\left(\beta_i^N, \sqrt{N}\sigma_{\frac{i-1}{N}}\Delta_{i+1}^N B\right) \right] + g\left(\beta_i^N, \sqrt{N}\sigma_{\frac{i-1}{N}}\Delta_{i+1}^N B\right)$$

so we may write

$$\mathbf{U}_{1N} = \frac{1}{N} \sum_{i=1}^{N-1} \mathbb{E}_{i-1} \left[g(\beta_i^N, \beta_{i+1}^N) - g\left(\beta_i^N, \sqrt{N}\sigma_{\frac{i-1}{N}}\Delta_{i+1}^N B\right) \right] + \frac{1}{N} \sum_{i=1}^{N-1} \sigma_{\frac{i-1}{N}}^2 \quad (17)$$

The first sum in (17) tends to zero in probability. To see this, note that the bound (9) implies the following limit in L_2 -norm:

$$\mathbb{E} \left| \frac{1}{N} \sum_{i=1}^{N-1} \mathbb{E}_{i-1} \left[g(\beta_i^N, \beta_{i+1}^N) - g\left(\beta_i^N, \sqrt{N}\sigma_{\frac{i-1}{N}}\Delta_{i+1}^N B\right) \right] \right|^2 \leq \frac{C}{N} \mathbb{E} \sum_{i=1}^{N-1} |\sigma_{\frac{i}{N}}^2 - \sigma_{\frac{i-1}{N}}^2|^2 \rightarrow 0 \quad (18)$$

where the convergence in (18) (which implies convergence in probability) follows from the fact that σ_t has finite quadratic variation. In addition, since $\{\sigma_t^2\}_{t \geq 0}$ is uniformly bounded and cadlag, the pointwise dominated convergence of $\sigma_u - \sigma_{\lfloor \frac{uN}{N} \rfloor} \rightarrow 0$ for $u \in [0; 1]$ follows, and Lebesgue's theorem yields

$$\sum_{i=1}^{N-1} \left[\int_{\frac{(i-1)}{N}}^{\frac{i}{N}} \left(\sigma_u^2 - \sigma_{\frac{(i-1)}{N}}^2 \right) du \right] \xrightarrow{a.s.} 0 \quad (19)$$

Together (18) and (19) imply $IV - U_{1N} \xrightarrow{P} 0$, which establishes (15). To show (16) we need the stronger assumption (A1). Define the sequence of independent standard normals $Z_i = \sqrt{N} \Delta_i^N B$, then Assumption (A1) yields

$$\mathbb{E}_{i-1} \left[\left(\sigma_{\frac{i}{N}}^2 - \sigma_{\frac{(i-1)}{N}}^2 \right) Z_{i+1}^2 \mathbf{1}_{Z_{i+1}^2 < Z_i^2} \right] = \mathbb{E}_{i-1} \left[\left(\sigma_{\frac{i}{N}}^2 - \sigma_{\frac{(i-1)}{N}}^2 \right) \varphi(Z_i^2) \right] = O_P(1/N) \quad (20)$$

since $\varphi(Z_i^2) = \mathbb{E}_i[Z_{i+1}^2 \mathbf{1}_{Z_{i+1}^2 < Z_i^2}]$ is an even function of the Brownian path $\{B_t\}_{(i-1)/N < t < i/N}$. Now the property (10) yields

$$\mathbb{E}_{i-1} \left[g(\beta_i^N, \beta_{i+1}^N) - g\left(\beta_i^N, \sqrt{N} \sigma_{\frac{i-1}{N}} \Delta_{i+1}^N B\right) \right] = O_P(1/N) \quad (21)$$

This ensures that the first term in (17) is asymptotically negligible, even when scaled up by \sqrt{N} . Hence, the remaining task is to show,

$$\sqrt{N} \left(\frac{1}{N} \sum_{i=1}^{N-1} \sigma_{\frac{i-1}{N}}^2 - IV \right) \xrightarrow{P} 0.$$

However, this is a common task in the proof of CLT for IV estimators and the method of proof is, by now, well established; see, e.g., BNGJPS where the result is shown for a general setting of which the current framework is a special case. A more intuitive and detailed exposition is provided by Barndorff-Nielsen, Graversen, Jacod, and Shephard (2006), henceforth BNGJS. ■

Lemma 5 *Under the maintained assumptions, we have*

$$U_{2N} \xrightarrow{P} 0 \quad (22)$$

Proof. To simplify notation, define the martingale difference sequence $\left\{ \frac{1}{N} \eta_i^N, \mathcal{F}_{\frac{i}{N}} \right\}_{i \geq 0}$:

$$\eta_i^N = g(\beta_i^N, \beta_{i+1}^N) - E_{i-1} [g(\beta_i^N, \beta_{i+1}^N)]$$

Note that $\mathbb{E}[(\eta_i^N)^2 | \mathcal{F}_{\frac{i-1}{N}}] \leq C$, so applying the Cauchy-Schwartz inequality,

$$\mathbf{V} \left[\frac{1}{N} \sum_{i=1}^N \eta_i^N \right] = \frac{1}{N} \mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N ((\eta_i^N)^2 + 2\eta_i^N \eta_{i+1}^N) \right] \leq \frac{C}{N} \mathbb{E} \left[\frac{1}{N} \sum_{i=1}^N \mathbb{E}[(\eta_i^N)^2 | \mathcal{F}_{\frac{i-1}{N}}] \right] \leq \frac{C}{N} \rightarrow 0.$$

The L_2 convergence implies $\frac{1}{N} \sum_{i=1}^N \eta_i^N \xrightarrow{P} 0$. ■

Lemma 6 *Under the maintained assumptions, we have,*

$$(\mathbf{V}_{1N} - \mathbf{U}_{1N}) \xrightarrow{P} 0. \quad (23)$$

Under Assumption (A1), we obtain,

$$\sqrt{N} (\mathbf{V}_{1N} - \mathbf{U}_{1N}) \xrightarrow{P} 0. \quad (24)$$

Proof. To establish (24), and thus also (23), we must show,

$$\sqrt{N} (V_{1N} - U_{1N}) = \frac{1}{\sqrt{N}} \sum_{i=1}^{N-1} \mathbb{E}_{i-1} \left[(g(\chi_i^N, \chi_{i+1}^N) - g(\beta_i^N, \beta_{i+1}^N)) \right] \rightarrow 0 \text{ as } N \rightarrow \infty. \quad (25)$$

Using the bound (9), it follows that,

$$\begin{aligned} \sqrt{N} (V_{1N} - U_{1N}) &\leq \frac{1}{\sqrt{N}} \mathbb{E} \left[\sum_{i=1}^{N-1} |g(\chi_i^N, \chi_{i+1}^N) - g(\beta_i^N, \beta_{i+1}^N)| \right] \leq \frac{C}{\sqrt{N}} \mathbb{E} \left[\sum_{i=1}^N |(\chi_i^N)^2 - (\beta_i^N)^2| \right] \\ &= \frac{C}{\sqrt{N}} \sum_{i=1}^N \left(\mathbb{E}_{i-1} \left[h(\sqrt{N} \Delta_i^N Y) - \sigma_{i-1}^2 \right] \right) \end{aligned}$$

where we have defined the function $h(x) = x^2$. This formulation maps directly into the setting of BNGJPS where the results of this lemma are proven in a more general setting and for a generic $h(x)$ function subject to regularity conditions. In particular, our h function trivially satisfies the continuous differentiability and polynomial growth conditions necessary for the applicability of their analysis. An accessible, albeit lengthy, account of the steps of the argument may be found in BNGJS (2006, pp. 713-719). So while this proof is quite involved, the above reformulation of the relevant inequalities, as they arise within our specific setting, allows us to simply refer to previously published work for the result. ■

Lemma 7 *Under the maintained assumptions, we have,*

$$(\mathbf{V}_{2N} - \mathbf{U}_{2N}) \xrightarrow{P} 0. \quad (26)$$

Moreover, we may strengthen this result further to obtain,

$$\sqrt{N} (\mathbf{V}_{2N} - \mathbf{U}_{2N}) \xrightarrow{P} 0. \quad (27)$$

Proof. In order to demonstrate the second result of the lemma, which obviously implies the first, we define,

$$\xi_i^N = (1/\sqrt{N}) \left[g(\chi_i^N, \chi_{i+1}^N) - g(\beta_i^N, \beta_{i+1}^N) \right],$$

and we must then prove that,

$$\sum_{i=1}^{N-1} (\xi_i^N - \mathbb{E}_{i-1}[\xi_i^N]) \xrightarrow{P} 0$$

This expression constitutes a martingale difference sequence with respect to the filtration $\mathcal{F}_{\frac{i}{N}}$, so it suffices to show,

$$\sum_{i=1}^{N-1} \mathbb{E} [(\xi_i^N)^2] = \mathbb{E} \left[\sum_{i=1}^{N-1} \mathbb{E}_{i-1} [(\xi_i^N)^2] \right] \rightarrow 0 \text{ as } N \rightarrow \infty$$

Mimicking the type of steps undertaken in the proof of the previous lemma, including application of the uniform bound on moments of χ_i^N and β_i^N , we obtain,

$$\begin{aligned} \sum_{i=1}^{N-1} \mathbb{E} [(\xi_i^N)^2] &= \frac{1}{N} \mathbb{E} \left[\sum_{i=1}^{N-1} \mathbb{E}_{i-1} \left| g(\chi_i^N, \chi_{i+1}^N) - g(\beta_i^N, \beta_{i+1}^N) \right|^2 \right] \\ &\leq \frac{C}{N} \mathbb{E} \left[\sum_{i=1}^N \mathbb{E}_{i-1} \left[(h(\chi_i^N) - h(\beta_i^N))^2 \right] \right]. \end{aligned}$$

As for the previous lemma, our reformulation of the task maps the problem into the corresponding task in BNGJPS (2006) who prove the current lemma in a more general setting. A detailed account of the requisite steps to complete the proof may again be gleaned from BNGJS (2006, pp. 704-706). ■

Taken together, Lemma 4 - 5 and the first parts of Lemma 6 - 7 imply the consistency of our estimator under the minimal maintained assumptions. The second part of Lemma 6 - 7 is critical for the proof of the central limit theorem below.

A.2 Proposition 2: The CLT

Lemma 8 *Under assumption (A1), we have*

$$\sqrt{N} \mathbf{U}_{2N}^M \xrightarrow{\text{stable } \mathcal{D}} N \left(0, \nu \int_0^1 \sigma^4 du \right) \quad (28)$$

where the constant $\nu = \text{Var}[g(U_0, U_1)] + 2 \text{Cov}[g(U_0, U_1), g(U_1, U_2)]$ for $U_0, U_1, U_2 \sim i.i.d.N(0, 1)$.

Proof. Consider splitting the N scaled return observations into K blocks, the k^{th} of which is the vector $\chi_k^M = \{\sqrt{N} \Delta_i^N Y\}_{i \in \{(k-1)M+1, \dots, kM\}}$. The corresponding vector of observations from the approximating Brownian motion where volatility is held constant over the block is $\beta_k^{N,M} = \{\beta_i^{N,M}\}_{i \in \{(k-1)M+1, \dots, kM\}}$. Next, define by $g_M(\cdot) : \mathbb{R}^M \mapsto \mathbb{R}$ the block estimator of volatility:

$$g_M(\beta_k^{N,M}) = \frac{1}{M} \sum_{i=(k-1)M+1}^{kM-1} g(\beta_i^{N,M}, \beta_{i+1}^{N,M}) \quad (29)$$

We wish to apply Theorem IX.7.28 in Jacod and Shiryaev (2003) to $\sqrt{N} \mathbf{U}_{2N}^M$. Defining the martingale difference sequence $\psi_k^{N,M} = \sqrt{M} \left(g_M(\beta_k^{N,M}) - \frac{M-1}{M} \sigma_{\frac{(k-1)}{K}}^2 \right)$ we can write

$$\begin{aligned} \sqrt{N} \mathbf{U}_{2N}^M &= \frac{1}{\sqrt{K}} \sum_{k=1}^K \psi_k^{N,M} + \frac{1}{\sqrt{N}} \sum_{k=1}^{K-1} \left(g(\beta_{kM}^{N,M}, \beta_{kM+1}^{N,M}) - \mathbb{E}_{\frac{k-1}{K}} [g(\beta_{kM}^{N,M}, \beta_{kM+1}^{N,M})] \right) \\ &= \frac{1}{\sqrt{K}} \sum_{k=1}^K \psi_k^{N,M} + o_P(1) \end{aligned} \quad (30)$$

The last equality follows from the fact that each term in the second sum is centered and has bounded variance (given the uniform bound on σ_t). Thus the sum divided by \sqrt{N} will tend to zero provided $K = o_P(N)$.

We must now verify conditions (7.27)-(7.31) of Theorem IX.7.28. First note that $E[\psi_k^{N,M} | \mathcal{F}_{\frac{k-1}{K}}] = 0$ so that condition (7.27) is trivially satisfied. Condition (7.28) follows from the fact that

$$\frac{1}{K} \sum_{k=1}^K E \left[\left\{ \sqrt{M} \left(g_M(\beta_k^M) - \frac{M-1}{M} \sigma_{\frac{(k-1)}{K}}^2 \right) \right\}^2 \middle| \mathcal{F}_{\frac{k-1}{K}} \right] = \frac{\nu}{K} \sum_{k=1}^K \sigma_{\frac{k-1}{K}}^4 \xrightarrow{P} \nu \int_0^1 \sigma_u^4 du \quad (31)$$

where the convergence in probability (and in fact a.s.) is a consequence of the volatility process being cadlag and uniformly bounded. Next, we turn to condition (7.29). Let $\Delta_k^M \mathbf{B} = \left(B_{\frac{k}{K}} - B_{\frac{(k-1)}{K}}\right)$, then $E \left[\psi_k^{N,M} \Delta_k^M \mathbf{B} \middle| \mathcal{F}_{\frac{k-1}{K}} \right] = 0$, which follows from the fact that the variables $\psi_k^{N,M}$ are centered and that g_M is an even function. Condition (7.30), stating that $E \left[(\psi_k^{N,M})^2 \mathbf{1}_{|\psi_k^{N,M}| > \varepsilon} \right] \xrightarrow{P} 0$, follows straightforwardly from the fact that σ is uniformly bounded.

Finally, let $\{N_t\}_{t \in [0;1]}$ be a bounded martingale orthogonal to B (i.e. the covariation $\langle B, N \rangle_t = 0$ a.s.). We want to show that, for each block k , $E[\psi_k^{N,M} \left(N_{\frac{k}{K}} - N_{\frac{(k-1)}{K}}\right) | \mathcal{F}_{\frac{k-1}{K}}] = 0$. For $t > \frac{k-1}{K}$ consider the martingale difference sequence $M_t = E[\psi_k^{N,M} | \mathcal{F}_t]$. By the martingale representation theorem, $M_t = M_{\frac{k-1}{K}} + \int_{\frac{k-1}{K}}^t \varphi_u dB_u$ for some predictable process φ_u . Therefore the processes $\{M_t\}_{t > \frac{k-1}{K}}$ and $\{N_t - N_{\frac{k-1}{K}}\}_{t > \frac{k-1}{K}}$ are orthogonal and the product, $\{M_t(N_t - N_{\frac{k-1}{K}})\}$ is again a martingale which must then have mean zero. This verifies condition (7.31). Theorem IX.7.28 in Jacod and Shiryaev (2003) then implies that as N (and hence K and M) tend to infinity:

$$\sqrt{N} U_{2N}^M \xrightarrow{\text{stable}} N \left(0, \nu \int_0^1 \sigma^4 du \right) \quad (32)$$

■

Lemma 9 *Under the maintained assumptions, we have*

$$\sqrt{N} (U_{2N} - U_{2N}^M) \xrightarrow{P} 0 \quad (33)$$

Proof. Defining $\eta_i^{N,M} = g(\beta_i^{N,M}, \beta_{i+1}^{N,M}) - \mathbb{E}_{[(i-1)/M]M} [g(\beta_i^{N,M}, \beta_{i+1}^{N,M})]$, we note that $\left\{ \frac{1}{\sqrt{N}} (\eta_i^N - \eta_i^{N,M}) \right\}_{i \geq 1}$ is a martingale difference sequence with respect to the filtration $\{\mathcal{F}_{i/N}\}$. To show that $\sqrt{N} (U_{2N} - U_{2N}^M) = \sum_{i=1}^{N-1} (\eta_i^N - \eta_i^{N,M}) / \sqrt{N} \rightarrow 0$ in probability, it therefore suffices (by Doob's inequality, e.g. Revuz and Yor (1999), p.54-55) to show that

$$\frac{1}{N} \mathbb{E} \left[\sum_{i=1}^{N-1} |g(\beta_i^N, \beta_{i+1}^N) - g(\beta_i^{N,M}, \beta_{i+1}^{N,M})|^2 \right] \rightarrow 0 \quad (34)$$

By the bound of $g(\cdot)$ we have

$$\begin{aligned} \frac{1}{N} \mathbb{E} \left[\sum_{i=1}^{N-1} |g(\beta_i^N, \beta_{i+1}^N) - g(\beta_i^{N,M}, \beta_{i+1}^{N,M})|^2 \right] &\leq \frac{C}{N} \mathbb{E} \left[\sum_{i=1}^N \mathbb{E}_{i-1} |(\beta_i^N)^2 - (\beta_i^{N,M})^2|^2 \right] \\ &\leq \frac{C}{N} \mathbb{E} \left[\sum_{i=1}^N \left| \sigma_{\frac{i-1}{N}}^2 - \sigma_{\frac{[(i-1)/M]M}{N}}^2 \right|^2 \right] = C \mathbb{E} \int_0^1 \left(\sigma_{\frac{[uN]}{N}}^2 - \sigma_{\frac{[uK]}{K}}^2 \right)^2 du = o_P(1) \end{aligned} \quad (35)$$

where the last inequality follows from the uniform boundedness of σ_t and Lebesgue's theorem. ■

Importantly, the specification of the volatility process in Assumption (A1) can be extended to include finite as well as infinite activity jump processes subject only to very weak regularity conditions, stipulating that the volatility process evolves according to an Ito semimartingale where the jump components have locally bounded jump characteristics, as laid out in BNGJPS. This follows from the fact that the only terms in (14) affected by the inclusion of jumps are $\sqrt{N}(V_{1N} - U_{1N})$ and $\sqrt{N}(V_{2N} - U_{2N})$ which map into the corresponding terms in BNGJPS as outlined in the proofs above. As such, the distributional results of the paper cover a very wide range of underlying return generating processes.

Finally, we note that the proof for the MedRV estimator follows analogously by simply changing the g function accordingly. The proof of Proposition 3 is omitted, but it may be derived using the identical strategy, in which volatility is held constant over blocks of increasing size. In particular, the conditional covariance between the estimators can easily be calculated on each block and a stable convergence argument similar to Lemma 8 goes through.

A.3 The Asymptotic Distribution under Jump Alternatives

Suppose now the log price process is given as $X = Y + J$, where Y is a Brownian semi-martingale of the form (6) while J is a finite activity jump process. While the specification of the jump process is restrictive, it covers many cases of interest and it can be generalized to infinite activity jump processes along the lines of Barndorff-Nielsen, Shephard, and Winkel (2006). Restating their Proposition 1 in our notation, we have $\sqrt{N}|Y_{\frac{i}{N}} - Y_{\frac{i-1}{N}}| = O_P(|\log(N)|^{1/2})$, which, as they show, follows from Levy's modulus of continuity theorem for Brownian motion. This immediately yields:

Proposition 10 *When J is a **finite activity** jump process, the asymptotic distribution of the MinRV and MedRV estimators applied to the processes $\{X_t\}$ and $\{Y_t\}$ are identical.*

Proof.

As before, we deal only with the MinRV case as the MedRV case is analogous. On a given realization of the path there is a finite number of jumps, so (asymptotically) at most one of the terms $|X_{\frac{i}{N}} - X_{\frac{i-1}{N}}|$ or $|X_{\frac{i+1}{N}} - X_{\frac{i}{N}}|$ will include a jump. Therefore, each term in the estimator (up to a normalizing constant) is

$$\min \left(|X_{\frac{i}{N}} - X_{\frac{i-1}{N}}|^2, |X_{\frac{i+1}{N}} - X_{\frac{i}{N}}|^2 \right) = O_P \left(\frac{\log N}{N} \right)$$

regardless of whether a (single) jump occurred or not. Since only finitely many terms differ,

$$\begin{aligned} \sum_{j=1}^N \left[\min \left(|X_{\frac{j}{N}} - X_{\frac{j-1}{N}}|^2, |X_{\frac{j+1}{N}} - X_{\frac{j}{N}}|^2 \right) - \min \left(|Y_{\frac{j}{N}} - Y_{\frac{j-1}{N}}|^2, |Y_{\frac{j+1}{N}} - Y_{\frac{j}{N}}|^2 \right) \right] \\ = O_P \left(\frac{\log N}{N} \right) = o_P \left(\frac{1}{\sqrt{N}} \right) \end{aligned}$$

so neither the consistency nor the convergence in distribution is affected by the occurrence of finite activity jumps.

■

B Figures and Tables




Figure 1: **Average IV estimates across 33 DJ30 stocks between January 1 2005 and May 31, 2007.** The first column (Panels (1A)-(3A)) shows the average IV estimates produced by each sub-sampled estimator as a function of sampling frequency (measured in seconds on the x-axis). Panel (1A) contains the calendar time estimates, Panel (2A) the tick time estimates, and Panel (3A) the tick time estimates after filtering out duplicate quotes. The second column (Panels (1B)-(3B)) plots the corresponding estimates for the pre-averaged version of each estimator as a function of pre-averaging window (measured in seconds on the x-axis), where the pre-averaging is carried out at the highest frequency as described in section 3.3.




Figure 2: **Average standard deviation of IV estimates across 33 DJ30 stocks between January 1 2005 and May 31, 2007.** The first column (Panels (1A)-(3A)) shows the average standard deviation of the IV estimates produced by each sub-sampled estimator as a function of sampling frequency (measured in seconds on the x-axis). Panel (1A) contains the calendar time estimates, Panel (2A) the tick time estimates, and Panel (3A) the tick time estimates after filtering out duplicate quotes. The second column (Panels (1B)-(3B)) plots the corresponding standard deviation for the pre-averaged version of each estimator as a function of pre-averaging window (measured in seconds on the x-axis), where the pre-averaging is carried out at the highest frequency as described in section 3.3.




Figure 3: Average IV estimates across 33 DJ30 stocks between June 1 2007 and May 31, 2008. The first column (Panels (1A)-(3A)) shows the average IV estimates produced by each sub-sampled estimator as a function of sampling frequency (measured in seconds on the x-axis). Panel (1A) contains the calendar time estimates, Panel (2A) the tick time estimates, and Panel (3A) the tick time estimates after filtering out duplicate quotes. The second column (Panels (1B)-(3B)) plots the corresponding estimates for the pre-averaged version of each estimator as a function of pre-averaging window (measured in seconds on the x-axis), where the pre-averaging is carried out at the highest frequency as described in section 3.3.




Figure 4: **Average standard deviation of IV estimates across 33 DJ30 stocks between June 1 2007 and May 31, 2008.** The first column (Panels (1A)-(3A)) shows the average standard deviation of the IV estimates produced by each sub-sampled estimator as a function of sampling frequency (measured in seconds on the x-axis). Panel (1A) contains the calendar time estimates, Panel (2A) the tick time estimates, and Panel (3A) the tick time estimates after filtering out duplicate quotes. The second column (Panels (1B)-(3B)) plots the corresponding standard deviation for the pre-averaged version of each estimator as a function of pre-averaging window (measured in seconds on the x-axis), where the pre-averaging is carried out at the highest frequency as described in section 3.3.

Table 1: **Variance factors for multipower and MinRV and MedRV estimators.** All estimators have an asymptotic variance of the form $\nu \int_0^1 \sigma_u^4 du$. The table displays the variance factor ν for each estimator.

	RV	BV	MedRV	MPV(3)	MPV(4)	MPV(5)	MPV(6)	MinRV	MPV(7)
Variance Factor	2.00	2.61	2.96	3.06	3.38	3.61	3.78	3.81	3.91

Table 2: Descriptive statistics for the 33 stocks that were part of the Dow Jones 30 index between January 1, 2005 and May 31, 2007.

Symbol	Average # Quotes	Average # Non-Duplicate Quotes	Average Price (\$)	Average 2-min RV ($\times 10^{-4}$)	Average 2-min BV ($\times 10^{-4}$)	Average Log Spread ($\times 10^{-4}$)	Average Ratio Log Spread/Sigma	Average # zero returns per day lasting above 30s / 60s / 120s
AA	11,062	3,610	30	2.22	2.10	4.80	0.03	161 / 48 / 11
AIG	11,420	4,484	64	1.00	0.95	2.73	0.03	127 / 30 / 5
AXP	10,280	3,444	55	0.81	0.76	2.84	0.03	161 / 48 / 11
BA	10,481	4,919	74	1.26	1.19	2.97	0.03	113 / 21 / 3
BAC	12,364	3,496	48	0.68	0.64	2.87	0.04	161 / 49 / 11
C	13,201	4,073	49	0.77	0.73	2.70	0.03	144 / 42 / 9
CAT	9,660	4,361	71	1.62	1.54	3.01	0.02	131 / 27 / 4
CVX	13,353	6,169	63	1.64	1.58	2.88	0.02	82 / 15 / 2
DD	10,460	3,597	45	1.19	1.12	3.48	0.03	154 / 45 / 10
DIS	11,421	2,774	29	1.08	1.01	4.46	0.04	177 / 67 / 19
EK	7,645	1,769	26	2.19	1.93	5.48	0.04	189 / 81 / 28
GE	14,010	2,565	35	0.63	0.59	3.35	0.04	182 / 70 / 20
GM	10,225	3,211	30	3.70	3.49	5.09	0.03	162 / 51 / 13
HD	11,606	3,814	39	1.35	1.28	3.69	0.03	149 / 39 / 8
HPQ	12,284	3,324	32	1.61	1.50	4.32	0.04	163 / 55 / 14
IBM	11,626	5,531	86	0.84	0.80	2.43	0.03	89 / 14 / 2
INTC	17,285	1,638	22	1.67	1.51	4.63	0.04	206 / 99 / 35
IP	8,064	2,277	34	1.36	1.25	4.28	0.04	191 / 71 / 20
JNJ	11,722	4,091	64	0.52	0.49	2.47	0.04	130 / 32 / 6
JPM	12,263	3,453	42	0.88	0.83	3.27	0.04	163 / 52 / 12
KO	11,269	3,158	44	0.59	0.55	3.15	0.04	166 / 57 / 15
MCD	11,171	3,238	36	1.22	1.13	3.90	0.04	164 / 55 / 14
MMM	9,421	4,026	78	0.91	0.87	2.74	0.03	141 / 31 / 5
MO	10,544	4,195	74	1.09	0.93	2.38	0.02	142 / 33 / 5
MRK	11,424	3,315	37	1.58	1.38	3.84	0.03	165 / 54 / 13
MSFT	16,575	1,272	27	0.94	0.84	3.80	0.04	199 / 106 / 43
PFE	13,654	2,501	26	1.12	1.06	4.63	0.05	184 / 71 / 21
PG	11,663	4,101	58	0.72	0.69	2.69	0.03	144 / 38 / 7
T	11,970	2,472	27	1.08	1.00	4.94	0.05	177 / 77 / 27
UTX	10,201	4,432	68	1.07	1.01	3.16	0.03	127 / 27 / 4
VZ	12,339	3,056	35	0.98	0.91	3.81	0.04	165 / 61 / 18
WMT	12,475	4,345	48	0.91	0.88	3.01	0.03	130 / 32 / 6
XOM	14,831	6,133	64	1.39	1.34	2.27	0.02	90 / 17 / 2
Mean ALL:	11,757	3,601	47	1.23090	1.14720	0.00035	0.03	152 / 49 / 13
Median ALL	11,778	3,601	48	1.08398	1.00552	0.00033	0.03	161 / 48 / 11
Max ALL:	17,285	6,169	86	3.69647	3.48752	0.00055	0.05	206 / 106 / 43
Min ALL:	7,645	1,272	22	0.52107	0.49326	0.00023	0.02	82 / 14 / 2

Table 3: **Relative bias and relative mean squared error (MSE) factors for sub-sampled IV estimators at the 12-second, 60-second, and 300-second sampling frequencies.** We report the bias and MSE factors for each sub-sampled estimator and each model based on sampling frequency 12 seconds (top panel), 60 seconds (middle panel), and 300 seconds (bottom panel). The MSE factor is computed as the sample mean of $390(\widehat{IV} - IV)^2/IQ$, where IV and IQ are the true simulated integrated variance and integrated quarticity on each day. The MSE factor for 60-sec sub-sampled RV is thus ≈ 1.33 which is the theoretical value shown by Zhang, Mykland, and Ait-Sahalia (2005). Each column corresponds to a given estimator: Realized volatility (RV), bipower variation (BV), tripower variation (TV), Realized Quantile RV (QRV), the MinRV and the MedRV. Each row corresponds to one of the models described in detail in Section 4.

	Relative Bias						Relative MSE					
	Sub-Sampled Estimators											
	RV	BV	TV	QRV	MinRV	MedRV	RV	BV	TV	QRV	MinRV	MedRV
12-sec frequency	Model 1: BM	1.000	1.000	0.999	1.000	1.000	0.268	0.310	0.337	0.300	0.389	0.328
	Model 2: SV-U	0.999	0.998	0.971	0.998	0.998	0.273	0.310	0.336	0.550	0.386	0.338
	Model 3: BM + Sparsity	0.999	0.983	0.977	0.978	0.971	0.345	0.476	0.580	0.525	0.772	0.618
	Model 4: BM + 1 Jump	1.244	1.021	1.011	1.002	1.002	75.196	0.636	0.402	0.295	0.384	0.337
	Model 5: BM + 4 Jumps	1.250	1.042	1.025	1.007	1.007	37.245	1.146	0.615	0.319	0.412	0.372
	Model 6: BM + Noise	1.083	1.084	1.084	1.082	1.084	2.949	3.041	3.061	2.964	3.148	3.059
60-sec frequency	Model 1: BM	1.000	1.000	0.999	1.000	0.999	1.350	1.511	1.613	1.455	1.857	1.633
	Model 2: SV-U	0.996	0.993	0.991	0.970	0.994	1.309	1.485	1.568	1.540	1.845	1.582
	Model 3: BM + Sparsity	1.000	0.996	0.995	0.993	0.994	1.436	1.586	1.698	1.528	1.952	1.710
	Model 4: BM + 1 Jump	1.242	1.044	1.027	1.008	1.008	75.595	3.135	2.199	1.562	2.006	1.753
	Model 5: BM + 4 Jumps	1.250	1.085	1.062	1.033	1.029	38.855	5.124	3.520	1.969	2.339	2.227
	Model 6: BM + Noise	1.017	1.017	1.017	1.017	1.018	1.454	1.681	1.769	1.546	2.065	1.798
300-sec frequency	Model 1: BM	1.001	1.002	1.002	1.002	1.002	6.736	7.808	8.342	7.394	9.676	8.432
	Model 2: SV-U	0.991	0.980	0.970	0.967	0.980	6.606	7.166	7.521	6.686	8.709	7.637
	Model 3: BM + Sparsity	1.002	1.002	1.003	1.000	1.002	6.913	7.891	8.436	7.447	9.783	8.588
	Model 4: BM + 1 Jump	1.242	1.089	1.066	1.039	1.031	81.665	15.012	12.275	8.462	10.704	9.672
	Model 5: BM + 4 Jumps	1.250	1.149	1.126	1.110	1.093	47.083	21.087	18.078	14.415	16.119	15.620
	Model 6: BM + Noise	1.003	1.005	1.005	1.004	1.007	6.852	7.753	8.289	7.406	9.599	8.246

Table 4: **Relative bias and relative mean squared error (MSE) factors for pre-averaged IV estimators using 12-second, 60-second, and 300-second pre-averaging windows.** We report the bias and MSE factors for each pre-averaged estimator and each model based on pre-averaging window of 12 seconds (top panel), 60 seconds (middle panel), and 300 seconds (bottom panel). The MSE factor is computed as the sample mean of $390(\widehat{IV} - IV)^2/IQ$, where IV and IQ are the true simulated integrated variance and integrated quarticity on each day. Each column corresponds to a given estimator: Realized volatility (RV), bipower variation (BV), tripower variation (TV), Realized Quantile RV (QRV), the MinRV and the MedRV. Each row corresponds to one of the models described in detail in Section 4.

	Relative Bias						Relative MSE					
	Pre-Averaged Estimators											
	RV	BV	TV	QRV	MinRV	MedRV	RV	BV	TV	QRV	MinRV	MedRV
12-sec window	Model 1: BM	1.000	1.000	1.000	0.999	1.000	1.000	0.251	0.276	0.232	0.334	0.271
	Model 2: SV-U	0.999	0.999	0.998	0.971	0.999	0.998	0.240	0.264	0.496	0.316	0.261
	Model 3: BM + Sparsity	0.999	0.974	0.965	0.969	0.955	0.962	0.311	0.601	0.680	1.195	0.907
	Model 4: BM + 1 Jump	1.244	1.018	1.009	1.001	1.002	1.002	75.350	0.479	0.222	0.329	0.270
	Model 5: BM + 4 Jumps	1.250	1.035	1.020	1.006	1.006	1.006	37.093	0.851	0.243	0.340	0.291
	Model 6: BM + Noise	1.078	1.079	1.079	1.078	1.079	1.079	2.616	2.683	2.718	2.784	2.715
60-sec window	Model 1: BM	1.001	1.000	1.000	1.000	1.000	1.000	1.288	1.415	1.202	1.679	1.386
	Model 2: SV-U	0.995	0.993	0.990	0.969	0.993	0.991	1.269	1.388	1.355	1.692	1.379
	Model 3: BM + Sparsity	1.001	0.993	0.991	0.988	0.988	0.990	1.197	1.380	1.313	1.825	1.512
	Model 4: BM + 1 Jump	1.242	1.038	1.023	1.006	1.007	1.007	75.497	2.536	1.247	1.826	1.515
	Model 5: BM + 4 Jumps	1.250	1.073	1.051	1.026	1.023	1.026	38.454	3.984	1.501	2.024	1.791
	Model 6: BM + Noise	1.003	1.004	1.003	1.003	1.004	1.004	1.090	1.333	1.456	1.761	1.430
300-sec window	Model 1: BM	1.001	1.001	1.002	1.001	1.002	1.002	5.374	6.654	6.002	9.004	7.321
	Model 2: SV-U	0.990	0.979	0.968	0.967	0.979	0.969	5.358	6.106	5.652	7.987	6.668
	Model 3: BM + Sparsity	1.002	1.001	1.003	0.998	1.000	1.002	5.569	6.793	6.241	9.220	7.546
	Model 4: BM + 1 Jump	1.241	1.075	1.053	1.031	1.024	1.027	80.465	11.892	9.993	9.608	8.055
	Model 5: BM + 4 Jumps	1.251	1.131	1.107	1.086	1.073	1.082	45.584	17.171	14.700	13.342	12.172
	Model 6: BM + Noise	1.002	1.004	1.004	1.001	1.005	1.002	5.655	6.711	7.285	8.794	7.255

- 2009-39: Borus Jungbacker, Siem Jan Koopman and Michel van der Wel: Dynamic Factor Models with Smooth Loadings for Analyzing the Term Structure of Interest Rates
- 2009-40: Niels Haldrup, Antonio Montañés and Andreu Sansó: Detection of additive outliers in seasonal time series
- 2009-41: Dennis Kristensen: Pseudo-Maximum Likelihood Estimation in Two Classes of Semiparametric Diffusion Models
- 2009-42: Ole Eiler Barndorff-Nielsen and Robert Stelzer: The multivariate supOU stochastic volatility model
- 2009-43: Lasse Bork, Hans Dewachter and Romain Houssa: Identification of Macroeconomic Factors in Large Panels
- 2009-44: Dennis Kristensen: Semiparametric Modelling and Estimation: A Selective Overview
- 2009-45: Kim Christensen, Silja Kinnebrock and Mark Podolskij: Pre-averaging estimators of the ex-post covariance matrix
- 2009-46: Matias D. Cattaneo, Richard K. Crump and Michael Jansson: Robust Data-Driven Inference for Density-Weighted Average Derivatives
- 2009-47: Mark Podolskij and Mathias Vetter: Understanding limit theorems for semimartingales: a short survey
- 2009-48: Isabel Casas and Irene Gijbels: Unstable volatility functions: the break preserving local linear estimator
- 2009-49: Torben G. Andersen and Viktor Todorov: Realized Volatility and Multipower Variation
- 2009-50: Robinson Kruse, Michael Frömmel, Lukas Menkhoff and Philipp Sibbertsen: What do we know about real exchange rate non-linearities?
- 2009-51: Tue Gørgens, Christopher L. Skeels and Allan H. Würtz: Efficient Estimation of Non-Linear Dynamic Panel Data Models with Application to Smooth Transition Models
- 2009-52: Torben G. Andersen, Dobrislav Dobrev and Ernst Schaumburg: Jump-Robust Volatility Estimation using Nearest Neighbor Truncation