



SCHOOL OF ECONOMICS AND MANAGEMENT
FACULTY OF SOCIAL SCIENCES
AARHUS UNIVERSITY



CREATES
Center for Research in Econometric
Analysis of Time Series

CREATES Research Paper 2009-54

Testing a parametric function against a nonparametric alternative in IV and GMM settings

Tue Gørgens and Allan Würtz

School of Economics and Management
Aarhus University
Bartholins Allé 10, Building 1322, DK-8000 Aarhus C
Denmark

Testing a parametric function against a nonparametric alternative in IV and GMM settings

Tue Gørgens* Allan Würtz[†]

November 2009

Abstract: This paper develops a specification test for functional form for models identified by moment restrictions, including IV and GMM settings. The general framework is one where the moment restrictions are specified as functions of data, a finite-dimensional parameter vector, and a nonparametric real function (an infinite-dimensional parameter vector). The null hypothesis is that the real function is parametric. The test is relatively easy to implement and its asymptotic distribution is known. The test performs well in simulation experiments.

Keywords: Generalized method of moments, specification test, nonparametric alternative, LM statistic, generalized arc-sine distribution.

JEL classification codes: C12, C14, C52.

Acknowledgment: We thank Joel Horowitz for sharing his computer code, Bo Honoré and Chris Skeels for comments on early drafts, and CREATES and the Danish Social Science Research Council for financial support.

*Address: SPEAR Centre, Research School of Social Sciences, Australian National University, Canberra ACT 0200, Australia. E-mail: tue.gorgens@anu.edu.au.

[†]Address: CREATES and School of Economics and Management, University of Aarhus, DK-8000 Aarhus C, Denmark. E-mail: awurtz@econ.au.dk.

1 Introduction

Generalized method of moments (GMM) and its special cases instrumental variables (IV) and two-stage least squares (2SLS) are frequently used to estimate parametric models in econometrics. These models specify moments as functions of data and a finite-dimensional parameter vector. The functional form is assumed to be known, apart from the parameters. In many applications, it is desirable to test the validity of the assumed functional form. In some cases there may be an obvious alternative model to test against. Often, however, there are no obvious alternatives. In this paper, we develop a test of functional form, which has power against models which specify the moments as functions of data, a finite-dimensional parameter vector, and a real function (an infinite-dimensional parameter vector).

Our test is based on the ideas of Aerts, Claeskens, and Hart (1999). They considered testing a parametric fit against a nonparametric alternative within several estimation frameworks: maximum likelihood, quasi-maximum likelihood, and general estimating equations. Their test is based on a sequence of LM test statistics, each designed to test against a specific parametric alternative. The sequence nests the null model, and in the limit it spans the class of models which can be written as functions of data, a finite-dimensional parameter vector and a real function. The LM statistics are divided by their degrees of freedom, and a single test statistic is constructed as the largest of these weighted LM statistics.

In this paper we extend these ideas to the testing of models which are formulated as restrictions on moment functions. Such models include regression models, models estimated by IV and, more generally, models estimated by GMM. In particular, our extension is applicable in overidentified models. There are two important new issues to consider when extending the original test to a GMM framework, namely identification of the model under the alternative and the selection of moment restrictions to use in the construction of the LM statistics. We discuss two approaches to the selection issue. For simplicity we shall refer to our extension as the GMM-ACH test.

Our test is relatively simple to implement. In particular, the asymptotic distribution

and hence the asymptotic critical values of the test are known. Moreover, since the test is based on LM statistics it is not necessary to estimate any alternative models, which is an advantage in some applications. We anticipate that in most applications performing the test involves, in principle, nothing more complicated than taking derivatives and inverting matrices.

There already exists specification tests for models that are formulated as restrictions on conditional moment functions (see Bierens, 1990; Whang, 2001; Donald, Imbens, and Newey, 2003; Tripathi and Kitamura, 2003; Horowitz, 2006). Horowitz's (2006) simulation evidence suggests that his test has significantly better power than the other tests. Implementing Horowitz's test can be nontrivial, in part because it is not asymptotically pivotal. This implies that the critical values must be computed specifically for each application.

We compare the performance of our test to some of the existing tests in a Monte Carlo study. The results confirm the finding in earlier papers regarding the performance of the existing tests, namely, that the test by Horowitz tends to have the better power. Our test, however, has power close to (and some cases better than) that of Horowitz's test in these simulations.

The paper is structured as follows. Section 2 introduces the GMM-ACH test in a simple IV setting. Section 3 contains the general extension to the GMM setting. We develop the test for the case where the infinite-dimensional parameter vector is an unknown function of a real variable. We briefly discuss the extension to functions of several variables in the concluding section. Section 4 presents examples. Section 5 concludes.

Throughout the paper, $0_{(a \times b)}$ denotes an $a \times b$ -dimensional matrix of 0s and $I_{(j)}$ denotes the j -dimensional identity matrix. The symbol 0 is also used to denote a function which maps the real line to the number 0.

2 A simple IV model

In this section, we use a simple IV setup to explain how the GMM-ACH test is constructed. In the first subsection, we consider a version of the GMM-ACH test which uses the

minimum number of moment restrictions required for each LM statistic. In the second subsection, we discuss a version which uses the same set of moment restrictions for all LM statistics. In the last subsection we present the results of a Monte Carlo study.

2.1 Minimum number of moment restrictions

The objective is to test a given parametric model against a nonparametric alternative model. Using subscript i to indicate a generic observation, let y_i be a scalar left-hand side variable, let x_i be a scalar right-hand side variable, and let z_i be a scalar instrument. Assume n independent observations are available. In this section the parametric model of interest is

$$y_i = x'_{0i}\beta^* + u_i, \quad \mathbf{E}(u_i|z_i) = 0, \quad \beta^* \in \mathbb{R}^2, \quad (1)$$

where $x_{0i} = (1, x_i)'$, β^* is an unknown two-dimensional parameter vector and u_i is an unobserved random variable. The nonparametric alternative model is the general nonlinear model given by

$$y_i = x'_{0i}\beta^* + \gamma^*(x_i) + u_i, \quad \mathbf{E}(u_i|z_i) = 0, \quad \beta^* \in \mathbb{R}^2, \quad \gamma^* \in \Gamma, \quad (2)$$

where $\gamma^* : \mathbb{R} \rightarrow \mathbb{R}$ is a function and Γ is a set of continuous functions. We assume that $0 \in \Gamma$, so that model (2) nests model (1). In terms of (2), the null hypothesis is that $\gamma^* = 0$ and the alternative hypothesis is that $\gamma^* \neq 0$.

The GMM-ACH test is based on four steps. The first step is to construct a sequence of nested parametric alternative models which approximate the nonparametric model (2). A series expansion of γ is used for this purpose. Let b_1, b_2, \dots be a sequence of basis functions ($b_k : \mathbb{R} \rightarrow \mathbb{R}$ for $k = 1, 2, \dots$) and assume that for each $\gamma \in \Gamma$ there are coefficients $\alpha_{\gamma 1}, \alpha_{\gamma 2}, \dots$ such that given any $\epsilon > 0$,

$$\mathbf{P}\left(\left|\gamma(x_i) - \sum_{k=1}^j \alpha_{\gamma k} b_k(x_i)\right| > \epsilon\right) \rightarrow 0 \quad \text{as } j \rightarrow \infty. \quad (3)$$

It is not difficult to satisfy (3). For example, if x_i is bounded and $b_k(x_i) = x_i^{k-1}$ for $k = 1, 2, \dots$ (a basis of power functions), then (3) follows from the Weierstrass theorem.¹ The sequence of nested parametric alternative models is based on the partial sums of the series expansion. Define the functions g_1, g_2, \dots by

$$g_j(x_i, \theta_j) = \sum_{k=1}^j \theta_{jk} b_k(x_i), \quad j = 1, 2, \dots, \quad (4)$$

where $\theta_j = (\theta_{j1}, \dots, \theta_{jj})'$ is a j -dimensional parameter vector. Condition (3) means that γ^* can be approximated arbitrarily well by $g_j(\cdot, \theta_j)$ by taking j large enough and choosing the appropriate θ_j . Let $\theta_1^*, \theta_2^*, \dots$ denote these “pseudo-true” parameter vectors. A sequence of approximate alternative models can therefore be constructed as²

$$y_i = x'_{0i} \beta^* + g_j(x_i, \theta_j^*) + u_i, \quad E(u_i | z_i) \simeq 0, \quad \beta^* \in \mathbb{R}^2, \quad \theta_j^* \in \mathbb{R}^j, \quad j = 1, 2, \dots \quad (5)$$

The notation $E(u_i | z_i) \simeq 0$ is short-hand for $E(u_i | z_i) \rightarrow 0$ as $j \rightarrow \infty$. In terms of (5), the null hypothesis is that $\theta_j^* = 0_{(j \times 1)}$ for all $j = 1, 2, \dots$ and the alternative hypothesis is that $\theta_j^* \neq 0_{(j \times 1)}$ for some $j = 1, 2, \dots$

The second step in the GMM-ACH test concerns the identification of the parameters in the null model and in the approximate alternative models. We have chosen to specify the models using the conditional moment restriction $E(u_i | z_i) = 0$. We assume that this conditional moment restriction identifies the parameters under the null as well as under the alternative. In practice, if x_i is continuously distributed, then it is convenient to base estimation and testing on unconditional moment restrictions. Since $E(u_i | z_i) = 0$ implies $E(u_i t(z_i)) = 0$ for any choice of (measurable) function $t : \mathbb{R} \rightarrow \mathbb{R}$, arbitrarily many unconditional moment restrictions can easily be constructed. At least 2 moment restrictions are needed to identify and estimate β^* , and at least $2 + j$ moment restrictions are needed to identify and test hypotheses about $(\beta^{*'}, \theta_j^{*'})'$. A natural choice of instrument

¹For an introduction to the use of series in econometrics, see for example Pagan and Ullah (1999).

²In practice, it may happen that x_{0i} and the basis functions used in the construction of g_j are collinear. Indeed, this happened in the power function basis example offered just above. Since we are not interested in the latter per se, the offending terms may simply be omitted from g_j .

for identifying the coefficient on $b_k(x_i)$ is $b_k(z_i)$.

We proceed here by constructing an GMM-ACH test based on using the minimum number of moment restrictions required in each calculation. In the next section we discuss a version of the test which uses the same set of moment restrictions in all calculations. Under the null, assume β^* in (1) is identified by the two unconditional moment restrictions,

$$\mathbf{E}(z_{0i}(y_i - x'_{0i}\beta^*)) = 0_{(2 \times 1)}, \quad \beta^* \in \mathbb{R}^2, \quad (6)$$

where $z_{0i} = (1, z_i)'$. Under the alternatives, assume that $(\beta^{*'}, \theta_j^{*'})'$ is identified by the $2 + j$ unconditional moment restrictions

$$\mathbf{E}(z_{ji}(y_i - x'_{0i}\beta^* - g_j(x_i, \theta_j^*))) \simeq 0_{(2+j \times 1)}, \quad \beta^* \in \mathbb{R}^2, \quad \theta_j^* \in \mathbb{R}^j, \quad j = 1, 2, \dots, \quad (7)$$

where $z_{ji} = (1, z_i, b_1(z_i), \dots, b_j(z_i))'$ for $j = 1, 2, \dots$

The third step in the GMM-ACH test is to calculate a statistic for testing the null against each of the approximate alternative models. There are several statistics which can be used to test model (1) against the models given in (5). Here we follow Aerts, Claeskens, and Hart (1999) and use LM statistics. First we estimate the model under the null by solving the empirical analogues of (6). That is, the estimator, $\tilde{\beta} = (n^{-1} \sum_{i=1}^n z_{0i}x'_{0i})^{-1} (n^{-1} \sum_{i=1}^n z_{0i}y_i)$, is the solution in β to

$$n^{-1} \sum_{i=1}^n z_{0i}(y_i - x'_{0i}\beta) = 0_{(2 \times 1)}. \quad (8)$$

Then we construct an LM test based on the fact that if the null is true, then the empirical analogues of (7) should be (approximately) satisfied when evaluated at the parameter estimate obtained under the null; that is, at $\beta = \tilde{\beta}$ and $\theta_j = 0_{(j \times 1)}$. Let M_j denote the empirical moments evaluated at the estimator obtained under the null; that is,³

$$M_j = n^{-1} \sum_{i=1}^n z_{ji}(y_i - x'_{0i}\tilde{\beta}), \quad j = 0, 1, \dots \quad (9)$$

³For simplicity the dependence of M_j (and other random matrices defined below) on n is suppressed in the notation.

Note that by construction the first two components of M_j are 0. For given j , the LM statistic has the form $R_j = M_j' \text{Var}(M_j)^- M_j$, where $\text{Var}(M_j)^-$ is a generalized inverse of the variance matrix of M_j or an estimate of that matrix. Define $x_{ji} = (1, x_i, b_1(x_i), \dots, b_j(x_i))'$ and the matrices

$$A_j = -n^{-1} \sum_{i=1}^n z_{ji} x'_{ji}, \quad j = 0, 1, \dots \quad (10)$$

and

$$B_j = \frac{1}{n} \left(n^{-1} \sum_{i=1}^n (y_i - x'_{0i} \tilde{\beta})^2 z_{ji} z'_{ji} \right), \quad j = 0, 1, \dots \quad (11)$$

Define also the matrices

$$H_j = \begin{bmatrix} 0_{(j \times 2)} & I_{(j)} \end{bmatrix}, \quad j = 1, 2, \dots \quad (12)$$

Then the LM test statistics can be defined as⁴

$$R_j = M_j' (A_j^{-1})' H_j' (H_j A_j^{-1} B_j (A_j^{-1})' H_j')^{-1} H_j A_j^{-1} M_j, \quad j = 1, 2, \dots \quad (13)$$

Given j , R_j has an asymptotic χ_j^2 -distribution under the null. For a discussion of this particular variant of the LM statistic, see Section 3 and Appendix A.

The fourth and final step in the GMM-ACH test is to construct an overall test statistic by taking the maximum over a sequence of weighted LM statistics. The weights are the reciprocal of the degrees of freedom of the individual statistics. Specifically, the GMM-ACH test statistic is

$$S_r = \max_{1 \leq j \leq r} (R_j/j), \quad (14)$$

⁴The simple IV setup with exact identification is almost a special case of the GEE setup considered by Aerts, Claeskens, and Hart (1999). Their LM statistic is valid only if A_j is symmetric (e.g. if $z_i = x_i$). They stated the LM statistic in a different form. Let $[X]_j$ denote the lower right $j \times j$ -submatrix of the $(2+j) \times (2+j)$ -matrix X or the last j elements of the $(2+j)$ -vector X , then R_j can be expressed as $R_j = [M_j]'_j [A_j^{-1}]_j ([A_j^{-1} B_j A_j^{-1}]_j)^{-1} [A_j^{-1}]_j [M_j]_j$.

where r is treated as a constant.⁵ In Section 3, we argue that the distribution of S_r under the null converges, as $r \rightarrow \infty$ and $n \rightarrow \infty$, to a distribution which does not depend on any unknown population characteristics. Hart (1997, p178) tabulated this distribution, and the 1%, 5% and 10% critical values are 6.75, 4.18 and 3.22. The requirement that $r \rightarrow \infty$ is not important; Aerts, Claeskens, and Hart (1999, p872) claimed that the asymptotic approximation is usually fine for critical values less than 10% as long as $r > 5$.

The weighting of the LM statistics means that the ordering of the terms in the series approximation matters for the numerical value of the GMM-ACH test statistic. This issue also arises in nonparametric estimation based on series. The advice from that literature is to ensure that “important terms” are at the beginning of the series (see e.g. Gallant, 1981).

2.2 Same set of moment restrictions

The version of the GMM-ACH test presented above is based on using the minimum number of moment restrictions required to identify the parameters under the null and the alternative hypotheses. The literature on hypothesis testing in IV and GMM settings (e.g. Engle, 1984; Newey and McFadden, 1994) usually recommends using the same set of moment restrictions under both the null and the alternative. In this case, (6) and (7) are replaced by

$$\mathbb{E}(z_{ri}(y_i - x'_{0i}\beta^*)) = 0_{(2+r \times 1)}, \quad \beta^* \in \mathbb{R}^2, \quad (15)$$

and

$$\mathbb{E}(z_{ri}(y_i - x'_{0i}\beta^* - g_j(x_i, \theta_j^*))) \simeq 0_{(2+r \times 1)}, \quad \beta^* \in \mathbb{R}^2, \quad \theta_j^* \in \mathbb{R}^j, \quad j = 1, \dots, r. \quad (16)$$

⁵In a likelihood framework, rejecting the null if S_r is large is equivalent to rejecting the null if the Akaike Information Criterion (AIC) of one of the alternative models is sufficiently larger than the AIC of the null model. For further discussion of the connection between the GMM-ACH and the AIC statistics, see Aerts, Claeskens, and Hart (1999).

Except in the case where $j = r$ there are more equations than unknown parameters in (15) and (16).

The 2SLS estimator, $\tilde{\beta}$, of β^* based on (15) is

$$\tilde{\beta} = (A'_0 W_r A_0)^{-1} A'_0 W_r \left(n^{-1} \sum_{i=1}^n z_{ri} y_i \right), \quad (17)$$

where $A_0 = -n^{-1} \sum_{i=1}^n z_{ri} x'_{0i}$, and the weight matrix is $W_r = (n^{-1} \sum_{i=1}^n z_{ri} z'_{ri})^{-1}$. For each j , LM statistics for testing $\theta_j^* = 0_{(j \times 1)}$ against $\theta_j^* \neq 0_{(j \times 1)}$ using (16) can be constructed as

$$R_j = M'_r W_r A_j J_j A'_j W_r M_r, \quad j = 1, 2, \dots \quad (18)$$

where

$$M_r = n^{-1} \sum_{i=1}^n z_{ri} (y_i - x'_{0i} \tilde{\beta}), \quad (19)$$

$$A_j = -n^{-1} \sum_{i=1}^n z_{ri} x'_{ji}, \quad j = 0, \dots, r, \quad (20)$$

$$B_r = \frac{1}{n} \left(n^{-1} \sum_{i=1}^n (y_i - x'_{0i} \tilde{\beta})^2 z_{ri} z'_{ri} \right), \quad (21)$$

$$C_j = (A'_j W_r A_j)^{-1} A'_j W_r B_r W_r A_j (A'_j W_r A_j)^{-1}, \quad j = 1, \dots, r, \quad (22)$$

$$J_j = (A'_j W_r A_j)^{-1} H'_j (H_j C_j H'_j)^{-1} H_j (A'_j W_r A_j)^{-1}, \quad j = 1, \dots, r, \quad (23)$$

and H_j is defined in (12). Finally, the GMM-ACH statistic is $S_r = \max_{1 \leq j \leq r} (R_j/j)$, as before.

The asymptotic distributions of the LM statistics and the GMM-ACH statistic are the same as in the previous section. Detailed arguments are given in Section 3 and Appendix A.

When the minimum number of moment restrictions are used, the LM statistics are large if the additional instruments in z_{ji} are correlated with the residuals from the null

model. The LM statistics do not depend on the last j components of x_{ji} . When the same set of moment restrictions are used, z_{ri} is fixed and there are no additional instruments. The LM statistics are large if the columns in A_j corresponding to the additional regressors in x_{ji} are not orthogonal to the weighted empirical moments from the null model, $W_r M_r$. Since the first depends on additional instruments and the other on additional regressors, the two versions of the test may have different power properties. The next subsection presents a small Monte Carlo study which compares the two versions of the GMM-ACH test.

2.3 A small Monte Carlo study

In the remainder of this section we present and discuss simulation results on the finite-sample behavior of several versions of the GMM-ACH test for the simple IV model. We consider both the test based on the minimum number and on the same set of moment restrictions, and we calculate the tests using both power and Fourier flexible form bases in the series approximation. We compare the GMM-ACH tests with the tests developed by Donald, Imbens, and Newey (2003) and Horowitz (2006), as well as with simple ad hoc t and LM tests.

The test by Donald, Imbens, and Newey (2003) is based on the well-known Sargan (Hansen) test for overidentifying restrictions. In general, the Sargan test does not have power against nonparametric alternatives. Donald, Imbens, and Newey modified the Sargan test by letting the number of overidentifying restrictions depend on the sample size. As the sample size increases, the test gains power against a larger set of alternatives. The additional moment restrictions are generated from a conditional moment restriction, as described in Section 2.1.

Horowitz (2006) developed a test based on estimating the difference between the parametric null model and the nonparametric alternative. He proved that the power of his test is arbitrarily close to 1 uniformly over a class of alternatives whose distance from the null hypothesis is of order $n^{-1/2}$.

Other specification tests in the context of GMM estimation have been developed by

Bierens (1990) and Tripathi and Kitamura (2003).⁶ These test have inferior power properties in the simulations conducted by Horowitz (2006), and for simplicity we do not report on them here.

As a benchmark, we report a simple t test based on the model obtained by adding one additional term to the null model. Since in most cases this alternative coincides with the data-generating process, we expect this t test to have very good power properties. In practice, the data-generating process is likely to be more complicated and we would then expect a t test to have less favorable power properties.

Finally, to illustrate the effect of taking the maximum of weighted LM test statistics against a sequence of parametric alternatives, we also report on the properties of an ordinary LM test against the largest (r th) parametric alternative.

The designs, and some of the results, are taken from Horowitz (2006). The data-generating process for all these experiments is

$$y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3 + u_i, \quad (24)$$

$$x_i = \Phi(\rho v_{1i} + (1 - \rho^2)^{1/2} v_{2i}), \quad (25)$$

$$z_i = \Phi(v_{1i}), \quad (26)$$

$$u_i = 0.2(\eta v_{2i} + (1 - \eta^2)^{1/2} v_{3i}), \quad (27)$$

where Φ denotes the standard normal distribution function, v_{1i} , v_{2i} and v_{3i} are independent standard normal random variables, and β_0 , β_1 , β_2 , β_3 , ρ and η are scalar parameters which vary across designs.

The results are shown in Table 1. Technical details of the implementation are given in the table notes. The results in the first part of the table show that the different versions of the GMM-ACH test have good level control. The only exception is the design where the GMM-ACH test is based on 2SLS and a power function basis. In that design the GMM-ACH test rejects too much and, perhaps surprisingly, so does the t test. In most other cases the level is correct within Monte Carlo sampling error (± 1.4 percentage point

⁶See also Whang (2001).

for a 5% test) or it is too low. This may cause lower power.

The second part of Table 1 shows that the GMM-ACH tests have powers comparable to Horowitz's test in these designs, and in some cases even better power. The test by Donald, Imbens, and Newey has significantly lower powers in most of the designs. Notice also that the idea of combining a sequence of LM test statistics into the GMM-ACH test generally has a positive effect on power. For many of the designs, there is a power loss of about 20 percentage points when doing a single LM test rather than doing the GMM-ACH test.

In sum, it appears that the GMM-ACH test has good properties. The level is well controlled, and the power is close to that of Horowitz's test and much better than Donald, Imbens, and Newey's test. A power basis seems to yield better power than a Fourier flexible form basis. However, this is not surprising given that the data generating process is polynomial. The simulations do not show a clear favorite between using the minimum number or the same set of moment restrictions in the GMM-ACH test. Finally, we note that the power of the GMM-ACH test is generally higher than the power of the ad hoc LM test.

3 A GMM-based specification test

The previous section presented the main ideas of the GMM-ACH test in the context of a simple linear IV model. In this section we develop the GMM-ACH test for a general nonlinear model identified by moment restrictions. Our framework includes many models of interest in economics such as system of equations models (typically estimated by two-stage least squares) and dynamic panel data models with fixed effects (typically estimated by GMM). When the parameters are overidentified, these models are not included in the frameworks discussed by Aerts, Claeskens, and Hart (1999).

Some econometric models are stated in terms of conditional and others in terms of unconditional moment restrictions. Ultimately the estimation of most models is based on unconditional moment restrictions, and we therefore specify the general model in terms of unconditional moment restrictions. As mentioned in Section 2.1, parameters in a

model identified by conditional moment restrictions can relatively easily be identified by unconditional moment restrictions derived from the conditional moment restrictions.⁷

3.1 The general setup

The setting is the following. Assume n independent observations are available for analysis. Let v_i be a generic random vector of data, let β^* be an unknown h -vector of parameters, and let $\gamma^* : \mathbb{R} \rightarrow \mathbb{R}$ be an unknown function. Let F be a known infinite-dimensional vector of functions of these three quantities. The econometric model is cast in terms of a vector of moment restrictions,

$$\mathbb{E}(F(v_i, \beta^*, \gamma^*)) = 0_{(\infty \times 1)}, \quad \beta^* \in \mathbb{R}^h, \quad \gamma^* \in \Gamma, \quad (28)$$

where again Γ is a set of continuous functions. We assume that $0 \in \Gamma$. We also assume that (28) identifies β^* and γ^* . In general it is not possible to identify a function (equivalent to an infinitely-dimensional parameter) such as γ^* from a finite set of moment restrictions, which is why we allow F to be infinitely-dimensional. In general, γ^* can be any continuous function. In terms of (28), the null hypothesis is that $\gamma^* = 0$. The alternative hypothesis is that $\gamma^* \neq 0$.

The range of null and alternative models which can be cast in the form of (28) is very wide. We provide a few examples in Section 4. The generality of (28) and the fact that we have made few assumptions about γ^* and how γ^* interacts with v_i and β^* are great strengths of the GMM-ACH approach. Often, γ^* will simply be a function of one of the components of v_i . In multiple-equation models, γ^* may be a function of a different component of v_i in each equation. In general, the argument of γ^* may be a function involving both v_i and β^* as in single-index models.

The GMM-ACH approach to testing the null against the nonparametric alternative is based on approximating the unknown γ^* with a sequence of nested parametric alternatives, g_1, g_2, \dots ; the construction of this sequence is explained in Section 2.1. Since only a finite

⁷Under certain conditions, a conditional moment restriction and a countable number of unconditional moment restrictions are equivalent, see e.g. Donald, Imbens, and Newey (2003, p58).

number of parameters are unknown under the null and the parametric alternatives, they may be identified from a finite set of moment restrictions. For $j = 1, 2, \dots$, let F_j denote the first l_j components of F . Under the null, assume without loss of generality that β^* is identified (possibly overidentified) by the l_0 moment restrictions

$$\mathbf{E}(F_0(v_i, \beta^*, 0)) = 0_{(l_0 \times 1)}, \quad \beta^* \in \mathbb{R}^h. \quad (29)$$

Under the parametric alternatives $j = 1, 2, \dots$, assume similarly that β^* and θ_j^* are identified (possibly overidentified) by the l_j moment restrictions⁸

$$\mathbf{E}(F_j(v_i, \beta^*, g_j(\cdot, \theta_j^*))) \simeq 0_{(l_j \times 1)}, \quad \beta^* \in \mathbb{R}^h, \quad \theta_j^* \in \mathbb{R}^j, \quad j = 1, 2, \dots \quad (30)$$

In terms of the parameters of the approximating models, the null hypothesis can be restated as $\theta_j^* = 0_{(j \times 1)}$ for all $j = 1, 2, \dots$, while the alternative hypothesis is that $\theta_j^* \neq 0_{(j \times 1)}$ for at least one of $j = 1, 2, \dots$

We now review GMM estimation and LM testing. For convenience, define $\delta_0 = \beta$ and $\delta_j = (\beta', \theta_j')'$ for $j = 1, 2, \dots$. Then define $f_0(\cdot, \delta_0) = F_0(\cdot, \beta, 0)$ and $f_j(\cdot, \delta_j) = F_j(\cdot, \beta, g_j(\cdot, \theta_j))$ for $j = 1, 2, \dots$. The GMM criterion functions, q_j , are

$$q_j(\delta_j) = (1/2)m_j(\delta_j)'W_jm_j(\delta_j), \quad j = 0, 1, \dots, \quad (31)$$

where W_j are some $l_j \times l_j$ symmetric weight matrices, and m_j are estimators of the moments $\mathbf{E}(f_j(v_i, \delta_j))$, as a function of δ_j , defined by

$$m_j(\delta_j) = n^{-1} \sum_{i=1}^n f_j(v_i, \delta_j), \quad j = 0, 1, \dots \quad (32)$$

For each $j = 0, 1, \dots$, the first-order condition for a minimum at $\tilde{\delta}_j$ is $\mathbf{D}q_j(\tilde{\delta}_j) = 0_{(h+j \times 1)}$.

⁸As in Section 2.1, $\theta_1^*, \theta_2^*, \dots$ denote “pseudo-true” values. The notation $\mathbf{E}(F_j(v_i, \beta^*, g_j(\cdot, \theta_j^*))) \simeq 0_{(l_j \times 1)}$ is short-hand for $\mathbf{E}(F_j(v_i, \beta^*, g_j(\cdot, \theta_j^*))) \rightarrow 0_{(l_j \times 1)}$ as $j \rightarrow \infty$.

The derivatives of q_j with respect to δ_j are

$$\mathbf{D}q_j(\delta_j) = a_j(\delta_j)'W_jm_j(\delta_j), \quad j = 0, 1, \dots, \quad (33)$$

where a_j are the gradients of m_j ,

$$a_j(\delta_j) = n^{-1} \sum_{i=1}^n \mathbf{D}f_j(v_i, \delta_j), \quad j = 0, 1, \dots \quad (34)$$

Here $\mathbf{D}f_j$ denotes the $l_j \times h + j$ -matrix of partial derivative functions of f_j with respect to δ_j . In many applications, the moment functions are linear in the parameters and the first-order conditions can be solved analytically for $\tilde{\delta}_j$.

Define the “pseudo-true” parameter vector $\delta_j^* = (\beta^{*'}, \theta_j^{*'})'$ and define the restricted estimator as $\tilde{\delta}_{0j} = (\tilde{\delta}_0', 0'_{(j \times 1)})'$ for $j = 1, 2, \dots$. Define the matrices

$$H_j = \begin{bmatrix} 0_{(j \times h)} & I_{(j)} \end{bmatrix}, \quad j = 1, 2, \dots \quad (35)$$

With this notation, the null hypothesis can then be expressed as $H_j\delta_j^* = 0_{(j \times 1)}$ for $j = 1, 2, \dots$, while the alternative is that $H_j\delta_j^* \neq 0_{(j \times 1)}$ for some $j = 1, 2, \dots$.

LM statistics are based on the fact that if the null is true, then the derivative of the GMM criterion function for model j should be close to $0_{(h+j \times 1)}$ when evaluated at $\tilde{\delta}_{0j}$. For each j , LM statistics for testing $H_j\delta_j^* = 0_{(j \times 1)}$ against $H_j\delta_j^* \neq 0_{(j \times 1)}$ have the form

$$R_j = \mathbf{D}q_j(\tilde{\delta}_{0j})' \text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j}))^- \mathbf{D}q_j(\tilde{\delta}_{0j}), \quad j = 1, 2, \dots, \quad (36)$$

where $\text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j}))^-$ is a generalized inverse of the variance matrix of the gradient $\mathbf{D}q_j(\tilde{\delta}_{0j})$ or an estimate of that matrix. Note that the rank of $\text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j}))$ is j . We discuss estimation of $\text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j}))$ and $\text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j}))^-$ below.

The GMM-ACH statistic, S_r , is the maximum of a sequence of weighted LM statistics for testing the null hypothesis against the alternatives in the sequence, where the weights

are the reciprocal of the statistic's degrees of freedom. Specifically,⁹

$$S_r = \max_{1 \leq j \leq r} (R_j/j), \quad (37)$$

where r is some appropriately large integer.

In the theorems below we describe two cases where S_r is asymptotically pivotal; that is, under the null its asymptotic distribution does not depend on any unknown population quantities. Specifically, the asymptotic distribution is a transformation of the generalized arc-sine distribution, namely

$$P(S_r \leq s) \rightarrow \exp\left(-\sum_{k=1}^{\infty} \frac{P(\chi_k^2 > ks)}{k}\right) \quad \text{as } r \rightarrow \infty \text{ and } n \rightarrow \infty, \quad (38)$$

where χ_k^2 has a chi-square distribution with k degrees of freedom. As mentioned in Section 2.1, asymptotic critical values have been tabulated by Hart (1997) and the asymptotic approximation is expected to be a good for critical values less than 10% as long as $r > 5$.

The nesting properties of the moment restrictions are important in the derivation of the asymptotic distribution of the LM statistics and the GMM-ACH statistic. In particular, the nesting properties are used to ensure that each LM statistic is asymptotically χ_j^2 -distributed and the differences between R_{j-1} and R_j for $j = 2, 3, \dots$ are asymptotically uncorrelated. For ease of reference, we state them as Assumption 1.

Assumption 1 *Let $l_0 \leq l_1 \leq \dots$. For $j = 1, 2, \dots$, the first l_{j-1} components of $f_j(v_i, \delta_j)$ equal $f_{j-1}(v_i, \delta_{j-1})$ for all (v_i, δ_j) such that $\delta_j = (\delta'_{j-1}, 0)'$, and the restricted estimator is $\tilde{\delta}_{0j} = (\tilde{\delta}'_0, 0'_{(j \times 1)})'$.*

Define $M_j = m_j(\tilde{\delta}_{0j})$ and $A_j = a_j(\tilde{\delta}_{0j})$. The main technical implications of Assumption 1 are that the upper l_0 -subvector of M_j equals M_0 and that the upper left $l_0 \times h$ -submatrix A_j equals A_0 .

The theorems below require that each LM statistic is asymptotically χ_j^2 -distributed under the null. Our setup is not quite standard, and we have been unsuccessful in finding

⁹While the LM statistic is convenient, alternatively one could base the GMM-ACH test on Wald or distance metric tests.

the necessary results in the literature. Standard treatments of LM statistics assume that W_j is an estimate of the optimal weight matrix and that the restricted estimator is obtained from minimizing Dq_j with respect to δ_j subject to the restrictions $H_j\delta_j^* = 0_{(j \times 1)}$ (see e.g. Newey and McFadden, 1994, Section 9). In the present case, the weight matrix, W_j , is arbitrary and the LM statistic is evaluated at $\tilde{\delta}_{0j}$, which is obtained from solving a different problem, namely the unrestricted minimization of Dq_0 with respect to δ_0 . Using Assumption 1, an estimator of $\text{Var}(Dq_j(\tilde{\delta}_{0j}))$ is derived in Appendix A.1.

3.2 Minimum number of moment restrictions

The first case we consider is where the number of moment restriction used under the null and each parametric alternative equals the number of parameters in the corresponding model. The simple IV model discussed in Section 2.1 is an example of such a setup. In this case, the parameters are exactly identified both under the null and approximate alternative hypotheses. Only a minimum number of moment restrictions are used in each step. When $l_j = h + j$ for all $j = 0, 1, \dots$, then $Dq_j(\tilde{\delta}_{0j})'\text{Var}(Dq_j(\tilde{\delta}_{0j}))^{-1}Dq_j(\tilde{\delta}_{0j})$ is the same as $M_j'\text{Var}(M_j)^{-1}M_j$. Define

$$B_j = \frac{1}{n} \left(n^{-1} \sum_{i=1}^n f_j(v_i, \tilde{\delta}_{0j}) f_j(v_i, \tilde{\delta}_{0j})' \right), \quad j = 0, 1, \dots, \quad (39)$$

$$C_j = A_j^{-1} B_j (A_j')^{-1}, \quad j = 1, 2, \dots, \quad (40)$$

and

$$J_j = (A_j^{-1})' H_j' (H_j C_j H_j')^{-1} H_j A_j^{-1}, \quad j = 1, 2, \dots, \quad (41)$$

then J_j is an estimator of $\text{Var}(M_j)^{-1}$. In Appendix A.2, we show that the LM statistics simplify to¹⁰

$$R_j = M_j' J_j M_j, \quad j = 1, 2, \dots \quad (42)$$

¹⁰This formula has the same form as the LM statistic based on the quasi-maximum likelihood estimator given in Theorem 3.5 in the article by White (1982).

Of course, this is the same formula as (13) used in Section 2.1. IV estimators are invariant to the choice of weight matrix, which therefore also drops out of the formula for the LM statistics.

Theorem 1 below states sufficient conditions for R_j to be asymptotically χ_j^2 -distributed and provides the corresponding asymptotic distribution of S_r .¹¹

Theorem 1 *Assumption 1 holds and technical regularity conditions are satisfied. For each $j = 1, 2, \dots$, suppose $l_j = h + j$ and A_j is invertible. For each $j = 1, 2, \dots$, suppose there exists a nonstochastic matrix, Σ_j , such that $n^{1/2}M_j \rightarrow^d N(0_{(h+j \times 1)}, \Sigma_j)$ and $\text{Var}(M_j) \rightarrow^p \Sigma_j$ as $n \rightarrow \infty$ and such that the first h rows and columns of Σ_j consist of 0s and the lower right $j \times j$ submatrix of Σ_j is positive definite. Then under the null the asymptotic distribution of S_r is given in (38).*

3.3 Same set of moment restrictions

The second case we consider is where the same set of moment restrictions and weight matrix are used in the null model as well as in each of the r alternatives. That is, $l_j = l_0$ and $W_j = W_0$ for all $j = 1, 2, \dots$. This is the case usually considered in the literature on hypothesis testing in IV and GMM settings (e.g. Engle, 1984; Newey and McFadden, 1994). The 2SLS setup in Section 2.2 provides an example. Using Assumption 1, we show in Appendix A.3 that J_j is an estimator of $\text{Var}(\text{D}q_j(\tilde{\delta}_{0j}))^-$, where

$$J_j = (A_j' W_r A_j)^{-1} H_j' (H_j C_j H_j')^{-1} H_j (A_j' W_r A_j)^{-1}, \quad j = 1, 2, \dots, \quad (43)$$

$$C_j = (A_j' W_r A_j)^{-1} A_j' W_r B_r W_r A_j (A_j' W_r A_j)^{-1}, \quad j = 1, 2, \dots, \quad (44)$$

¹¹For simplicity we do not spell out the standard regularity conditions required for Taylor expansions to be valid, central limit theorems to hold, etc. As indicated in (38), the limiting distribution is valid for $r \rightarrow \infty$ as $n \rightarrow \infty$. To bound the behavior of the test statistic as $r \rightarrow \infty$, it is assumed that, for given $\pi > 1$ and for every $\epsilon > 0$, there is a positive integer j_0 such that $P(\max_{j_0 \leq j \leq r} R_j/j \leq (\pi + 1)/2) < \epsilon$ for all sufficiently large n . Here π denotes the critical value used in the test.

and B_j is defined in (39). The LM statistics simplify to¹²

$$R_j = M'_r W_r A_j J_j A'_j W_r M_r, \quad j = 1, 2, \dots \quad (45)$$

The theorem below is the equivalent of Theorem 1.

Theorem 2 *Assumption 1 holds and technical regularity conditions are satisfied. For each $j = 1, 2, \dots$, suppose $l_j = l_0$ and $W_j = W_0$. For each $j = 1, 2, \dots$, suppose there exists a nonstochastic matrix, Σ_j , such that $n^{1/2}Dq_j(\tilde{\delta}_{0j}) \rightarrow^d N(0_{(h+j \times 1)}, \Sigma_j)$ and $\text{Var}(Dq_j(\tilde{\delta}_{0j})) \rightarrow^p \Sigma_j$ as $n \rightarrow \infty$ and such that the first h rows and columns of Σ_j consist of 0s and the lower right $j \times j$ submatrix of Σ_j is positive definite. Then under the null the asymptotic distribution of S_r is given in (38).*

The proofs of the theorems are omitted, since they are similar to the proof of Theorem 3 by Aerts, Claeskens, and Hart (1999).

3.4 Remarks

We conclude this section with some remarks. First, because of the LM approach, parameter estimates need only be calculated once. In some applications, not having to estimate the model under the alternative is an advantage. For example, it is often difficult to estimate models when the first-order conditions are nonlinear in the parameters.

Second, note that essentially the same assumptions underpin both Theorem 2 and Theorem 1. In practice, one therefore has a choice of whether to implement the test using the same set of moment restrictions in each step or using the minimum number of moment restrictions.

Third, when the same set of moment restrictions and the same weight matrix are used under the null as well as under the parametric alternatives, then the estimator, with 0s appended as appropriate, computed by solving the unrestricted problem of minimizing $q_0(\delta_0)$ with respect to δ_0 is identical to the estimators obtained by solving the restricted

¹²If an optimal weight matrix is used, so W_r and B_r^{-1} are equivalent, then R_j in (18) is the same as LM_{2n} in Table 2 in the article by Newey and McFadden (1994). In this case J_j simplifies to $J_j = (A'_j W_r A_j)^{-1} H'_j (H_j (A'_j W_r A_j)^{-1} H'_j)^{-1} H_j (A'_j W_r A_j)^{-1}$ for $j = 1, 2, \dots$

problem of minimizing $q_j(\delta_j)$ with respect to δ_j subject to $H_j\delta_j = 0_{(j \times 1)}$ for $j = 1, 2, \dots$. This can be seen from inspecting the first-order conditions.

Fourth, it is possible that there are other cases where S_r is asymptotically pivotal. A key property of the LM statistics under Theorem 2 is that the first $j - 1$ components of $Dq_j(\tilde{\delta}_{0j})$ equal $Dq_{j-1}(\tilde{\delta}_{0,j-1})$ for all $j = 1, \dots, r$. Mathematically, there are ways of achieving this which do not require using the same set of moment restrictions for all $j = 0, 1, \dots$. Examining the first-order conditions, (33), reveals that the key property is also satisfied if the partial derivatives of the last $l_j - l_{j-1}$ components of the empirical moment function with respect to the first $j - 1$ components of the parameter vector are all 0 and the weight matrix is block-diagonal with 0s in the first l_{j-1} rows (columns) of the last $l_j - l_{j-1}$ last columns (rows). The first requirement means that the additional $l_j - l_{j-1}$ moment restrictions must not depend on the previous $j - 1$ parameters. If the moment restrictions are constructed by multiplying instruments and “residuals”, then the additional $l_j - l_{j-1}$ instruments must be orthogonal to the partial derivatives of the residuals with respect to the previous $j - 1$ parameters.¹³ Thus, while it may be possible to construct other LM-based GMM-ACH test statistics, the requirements are complicated and seem less generalizable. Hence, we do not further pursue this possibility.

4 Examples

In this section we consider two GMM-ACH tests for a linear model with endogenous right-hand side variables. We focus on testing the specification of the conditional mean function, because we believe this is the testing problem most often faced by applied researchers.

Consider a linear model where one or more of regressors are endogenous. Let y_i be a scalar random variable as in Section 2, but now let x_i and z_i be random vectors. Also, partition $x_i = (w_{1i}, w'_{2i})'$ where w_{1i} is scalar. A constant may be included in x_i and z_i .

¹³If the weight matrix is constructed using the second moments of a set of instruments, then the additional $l_j - l_{j-1}$ instruments must also be orthogonal to the previous $j - 1$ instruments.

Suppose the parametric model of interest is

$$y_i = w_{1i}\beta_1^* + w_{2i}'\beta_2^* + u_i, \quad \mathbf{E}(u_i|z_i) = 0, \quad \beta^* \in \mathbb{R}^h, \quad (46)$$

where $\beta^* = (\beta_1^*, \beta_2^{*'})'$ is an unknown parameter vector and u_i is an unobserved random variable. As before, assume that n independent observations are available.

Equations of this form arises often in economics. For example, let (46) represent an Engel curve where y_i is the share of total expenditure spent on certain items in household i , w_{1i} is the log of total expenditure on nondurables (as an indicator of permanent income), w_{2i} represents household characteristics, and z_i includes the variables in w_{2i} as well as household income as the instrument for total expenditure. Then this is the well-known Working-Leser specification of the Engel curve relationship.

As another example, consider a simultaneous equation system representing demand and supply of a certain good. Let y_i be the log of the total (equilibrium) quantity of the good traded in market i , let w_{1i} be the log of the (equilibrium) price of the good, let w_{2i} represent the characteristics of buyers in market i , and let z_i include the variables in w_{2i} as well as characteristics of suppliers. Then (46) represents the structural demand equation.

4.1 Nonlinear effect of a single regressor

The first alternative specification we consider allows for a nonlinear effect in w_{1i} . In the Engel curve example, the alternative model represents a nonlinear permanent income effect. In the market demand example, the alternative model allows for a nonlinear price elasticity. Formally, the nonparametric alternative model is

$$y_i = w_{1i}\beta_1^* + w_{2i}'\beta_2^* + \gamma^*(w_{1i}) + u_i, \quad \mathbf{E}(u_i|z_i) = 0, \quad \beta^* \in \mathbb{R}^h, \quad \gamma^* \in \Gamma, \quad (47)$$

where again Γ is a set of continuous functions with $0 \in \Gamma$. The approximate alternative models are

$$y_i = w_{1i}\beta_1^* + w'_{2i}\beta_2^* + g_j(w_{1i}, \theta_j^*) + u_i, \quad \mathbb{E}(u_i|z_i) \simeq 0, \quad \beta^* \in \mathbb{R}^h, \quad \theta_j^* \in \mathbb{R}^j, \\ j = 1, 2, \dots, \quad (48)$$

where g_1, g_2, \dots are series approximations of γ and $\theta_1^*, \theta_2^*, \dots$ are pseudo-true values as defined earlier.

The main issue in applying the GMM-ACH test is to choose moment restrictions to estimate β^* under the null and to identify $\theta_1^*, \theta_2^*, \dots$ under the alternative. There are many potential restrictions to choose from in this model, since the conditional moment restriction implies an infinite number of unconditional moment restrictions which can be used for estimation and testing. In practice, under the null, the model is virtually always estimated using the restrictions

$$\mathbb{E}(z_i(y_i - w_{1i}\beta_1^* - w'_{2i}\beta_2^*)) = 0_{(l_0 \times 1)}, \quad \beta^* \in \mathbb{R}^h, \quad (49)$$

where l_0 is the dimension of z_i . Section 3 shows that there are two ways to proceed under the alternative.

If the number of instruments is equal to the number of endogenous variables and the null model is exactly identified, one has the option to base the test on Theorem 1. Define $z_{0i} = z_i$ and $z_{ji} = (z_i, b_1(z_i^1), \dots, b_j(z_i^1))'$ for $j = 1, 2, \dots$, where z_i^1 denotes one of the instruments and b_1, b_2, \dots are the basis functions used in the series approximations. If w_{1i} is exogenous, the natural choice for z_i^1 is w_{1i} itself. If w_{1i} is endogenous, the natural choice is one of the variables excluded from x_i .¹⁴ The moment restrictions are then

$$\mathbb{E}(z_{ji}(y_i - w_{1i}\beta_1^* - w'_{2i}\beta_2^* - g_j(w_{1i}, \theta_j^*))) \simeq 0_{(l_j \times 1)}, \quad \beta^* \in \mathbb{R}^h, \quad \theta_j^* \in \mathbb{R}^j, \\ j = 1, 2, \dots, \quad (50)$$

¹⁴It is possible to derive optimal instruments when the unconditional moment restrictions are based on a conditional moment restriction, see e.g. Newey and McFadden (1994, Sections 5.3–5.4).

where the number of moment restrictions is $l_j = h + j$ for $j = 0, 1, \dots$

Define $x_{0i} = (w_{1i}, w'_{2i})'$ and $x_{ji} = (w_{1i}, w'_{2i}, b_1(w_{1i}), \dots, b_j(w_{1i}))'$ for $j = 1, 2, \dots$. Formally the matrices which are used in the LM statistics and the GMM-ACH test statistic are exactly as given in (9)–(13) in Section 2.1, with the symbols x_{ji} and z_{ji} as defined in the present section and with $\tilde{\beta}$ being the usual IV estimator. To base this test on Theorem 2 instead of Theorem 1, simply use formulae (17)–(23) in Section 2.2.

If the null model is overidentified, it is most natural to base testing on Theorem 2. In this case, the need to choose which moment restrictions to use to identify $\theta_1^*, \theta_2^*, \dots$ is perhaps even more apparent. At one extreme one can use powers of a single variable as in the previous case. At the other extreme one can use power of all instruments. In the latter case, z_{ri} is redefined as $z_{ri} = (z_i, b_1(z_i^1), \dots, b_r(z_i^1), \dots, b_1(z_i^p), \dots, b_r(z_i^p))'$, where z^1, \dots, z^p are the nonconstant elements of z_i . In either case, the test is calculated using formulae (17)–(23).

4.2 Nonlinear effect of an index

The second alternative specification we consider allows for a nonlinear effect in the index $w'_{2i}\beta^*$. Specifically,

$$y_i = w_{1i}\beta_1^* + w'_{2i}\beta_2^* + \gamma^*(w'_{2i}\beta^*) + u_i, \quad \mathbb{E}(u_i|z_i) = 0, \quad \beta^* \in \mathbb{R}^h, \quad \gamma^* \in \Gamma. \quad (51)$$

This alternative is related to the well-known RESET test for functional form. In the Engel curve and the market demand examples, one might consider this alternative in order to check the robustness of $\tilde{\beta}_1$ to misspecification of the influence of household characteristics or buyer characteristics.

The approximate alternative models are

$$y_i = w_{1i}\beta_1^* + w'_{2i}\beta_2^* + g_j(w'_{2i}\beta^*, \theta_j^*) + u_i, \quad \mathbb{E}(u_i|z_i) \simeq 0, \quad \beta^* \in \mathbb{R}^h, \quad \theta_j^* \in \mathbb{R}^j, \\ j = 1, 2, \dots \quad (52)$$

Since there are no obvious single candidate instruments for the index, Theorem 2 is better

suited than Theorem 1. Let x_{ji} and z_{ji} be as discussed in the last paragraph of Section 4.1. The test statistics based on Theorem 2 is given in (17)–(23), except for the A_j matrices which for $j > 0$ become

$$A_j = -n^{-1} \sum_{i=1}^n z_{ri} \begin{bmatrix} x'_{0i} & D_1 g_j(w'_{2i} \tilde{\beta}, 0_{(j \times 1)}) \end{bmatrix}, \quad j = 1, \dots, \quad (53)$$

where $D_1 g_j$ denotes the partial derivative of g_j with respect to its first argument. If a power basis is used, then $D_1 g_j(w'_{2i} \tilde{\beta}, 0_{(j \times 1)}) = ((w'_{2i} \tilde{\beta})^2, \dots, (w'_{2i} \tilde{\beta})^{1+j})$.

4.3 Empirical example

In this subsection, we apply the GMM-ACH test to the Engel curve model described earlier. We use the same data as Blundell, Duncan, and Pendakur (1998); BDP henceforth.¹⁵ The data come from the 1980–1982 British Family Expenditure Survey. The extract is limited to married or cohabiting couples with one or two children, living in Greater London or south-east England, where the head of the household is currently employed. For further details about the sample, including summary statistics, please see BDP’s article.

One of the models considered by BDP has the form (46). In our notation, y_i is the share of total expenditure spent on certain items, w_{1i} is log of total expenditure, w_{2i} is a dummy for having two instead of one child in the family, and z_i includes w_{2i} as well as total disposable income.

The alternative specification is given in (47). (Since w_{2i} is a dummy, the alternative given in (51) is not relevant.) Table 2 shows estimation results using different parametric specifications and different estimation methods. The OLS estimates are similar to those reported in Tables II–VII by BDP, although not identical. The GMM-ACH tests reject the linear specification for fuel, transport and (marginally) for other goods. To help understand the outcome of the GMM-ACH tests, the last panel of Table 2 shows IV estimates for a model which is quadratic in the log of total expenditure. The statistical significance of the t -statistics for the coefficients on the squared terms agree with the GMM-ACH tests in all cases (the marginal case of other goods is only significant at the

¹⁵These data are available from the Journal of Applied Econometrics’ data archive.

5.7% level).

BDP also tested the linear model against a nonparametric alternative. Their approach is much more complicated than ours and involves estimating the model under the nonparametric alternative, a notoriously difficult problem. Their conclusions are different from ours. They rejected the linear specification for alcohol and other goods and no other categories. While the differences in conclusions are interesting, further investigation beyond the scope of the present paper.

5 Concluding remarks

Inspired by Aerts, Claeskens, and Hart (1999), we suggest an GMM-ACH specification test of a parametric function against a nonparametric alternative. The test is developed for models which are identified by moment restrictions. The test requires only estimation under the null, and hence nonparametric estimation is not involved. The GMM-ACH test is asymptotically pivotal, which makes it easy to obtain critical values.

In a small Monte Carlo study, the GMM-ACH test has good level and power properties compared to existing tests. The test developed by Horowitz (2006) appears to have the best power of all, but it is difficult to perform. The GMM-ACH test has power that is close to that of Horowitz's test, and it is easy to carry out. The simulations also show that the GMM-ACH test has substantially higher power than an LM test of the null against a single, high-order parametric alternative. Hence, the idea of combination of test statistics against a sequence of parametric alternatives proves to be valuable.

We have focused on testing against a nonparametric function of a single variable. In some applications, it would be useful to be able to test against a function of several variables (or perhaps against several functions). Based on the analysis by Aerts, Claeskens, and Hart (2000), we anticipate that the main issues regarding this extension are related to its practical implementation, since the basis functions will be functions of several variables. The asymptotic theory presented in the present paper should remain valid.

Originally, our interest in testing for functional form in GMM settings was motivated by dynamic panel data models with fixed effects. This particular application is relatively

complex, partly because these models have several equations per subject and each equation has its own set of instruments, and partly because GMM estimation of these models in practice is often troubled by weak instruments. We intend to publish our results for this case separately.

A Estimating the variance of the GMM gradient

In this appendix, we derive the estimators of $\text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j}))^-$ given in Section 3. Section A.1 shows that $\text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j}))$ can be estimated consistently. Section A.2 shows that the LM statistic (36) simplifies to (42) in the case where $l_j = h + j$ for all $j = 1, 2, \dots$. Section A.3 establishes (43) for the case where $l_j = l_0$ for all $j = 1, 2, \dots$. Throughout this appendix j is a fixed integer.

Our arguments in Section A.1 are similar to those given by Newey and McFadden (1994, Section 9). The main differences are that we consider the case where the restricted estimator may be based on a subset of the moment restrictions and where the weight matrix, W_j , is arbitrary. Newey and McFadden considered the case where the moment restrictions are identical under the null and the alternative and where W_j is an estimate of the optimal weight matrix.

A.1 The general case

In this section, we show that $\text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j}))$, which appears in (36) in Section 3, can be estimated as

$$\text{Var}(\mathbf{D}q_j(\tilde{\delta}_{0j})) = T_j B_j T_j', \quad j = 0, 1, \dots, \quad (54)$$

where B_j is given in (39) and T_j are the matrices defined by

$$T_j = A_j' W_j [I_{(l_j)} - A_j N_{1j} (A_0' W_0 A_0)^{-1} A_0' W_0 N_{2j}], \quad j = 0, 1, \dots, \quad (55)$$

with

$$N_{1j} = \begin{bmatrix} I_{(h)} \\ 0_{(j \times h)} \end{bmatrix}, \quad j = 0, 1, \dots, \quad (56)$$

and

$$N_{2j} = \begin{bmatrix} I_{(l_0)} & 0_{(l_0 \times l_j - l_0)} \end{bmatrix}, \quad j = 0, 1, \dots. \quad (57)$$

We use two key properties of the testing problem set up in Section 3, namely that the restricted estimator is $\tilde{\delta}_{0j} = (\tilde{\delta}'_0, 0'_{(j \times 1)})'$ where $\tilde{\delta}_0$ is the solution to the unrestricted minimization problem $Dq_0(\tilde{\delta}_0) = 0_{(h \times 1)}$, and that the first l_0 components of $m_j(\tilde{\delta}_{0j})$ equal $m_0(\tilde{\delta}_0)$. These properties are implied by Assumption 1.

In general, the GMM gradient evaluated at the unrestricted estimator is identically equal to $0_{(h+j \times 1)}$. However, this is not the case when evaluated at the restricted estimator. A Taylor series expansion of $m_j(\tilde{\delta}_{0j})$ around the “pseudo-true” value, δ_j^* , implies

$$n^{1/2}Dq_j(\tilde{\delta}_{0j}) = a_j(\tilde{\delta}_{0j})'W_j[n^{1/2}m_j(\delta_j^*) + a_j(\tilde{\delta}_{0j})n^{1/2}(\tilde{\delta}_{0j} - \delta_j^*)] + o_p(1), \quad (58)$$

where Dq_j , m_j and a_j are defined in (31), (32) and (34) and W_j is a given weight matrix. Under standard regularity conditions, $a_j(\tilde{\delta}_{0j})$ and W_j converge in probability to matrices of (finite) constants. Therefore the main sources of variation for $n^{1/2}Dq_j(\tilde{\delta}_{0j})$ are the empirical moments, $n^{1/2}m_j(\delta_j^*)$, and the estimated parameters, $n^{1/2}(\tilde{\delta}_{0j} - \delta_j^*)$.

In the present context, the restricted estimator has the form $\tilde{\delta}_{0j} = (\tilde{\delta}'_0, 0'_{(j \times 1)})'$, where $\tilde{\delta}_0$ is the solution to the (unrestricted) estimation problem, $Dq_0(\tilde{\delta}_0) = 0_{(h \times 1)}$. Under the null, the “pseudo-true” value can be similarly partitioned, $\delta_j^* = (\delta_0^{*'}, 0'_{(j \times 1)})'$. To derive the distribution of $n^{1/2}(\tilde{\delta}_0 - \delta_0^*)$, note that a Taylor series expansion similar to (58) yields

$$n^{1/2}Dq_0(\tilde{\delta}_0) = a_0(\tilde{\delta}_0)'W_0[n^{1/2}m_0(\delta_0^*) + a_0(\tilde{\delta}_0)n^{1/2}(\tilde{\delta}_0 - \delta_0^*)] + o_p(1). \quad (59)$$

Since $Dq_0(\tilde{\delta}_0) = 0_{(h \times 1)}$, it follows that

$$n^{1/2}(\tilde{\delta}_0 - \delta_0^*) = -(a_0(\tilde{\delta}_0)'W_0a_0(\tilde{\delta}_0))^{-1}a_0(\tilde{\delta}_0)'W_0n^{1/2}m_0(\delta_0^*) + o_p(1). \quad (60)$$

An approximation for $n^{1/2}(\tilde{\delta}_{0j} - \delta_j^*)$ follows by appending rows of zeros. With N_{1j} as defined in (56), $n^{1/2}(\tilde{\delta}_{0j} - \delta_j^*) = N_{1j}n^{1/2}(\tilde{\delta}_0 - \delta_0^*)$. Before inserting into (58), it is convenient to express $m_0(\delta_0^*)$ in terms of $m_j(\delta_j^*)$. This will facilitate keeping track of the covariance between the empirical moments and the estimated parameters in (58). By construction, estimation under the null is based on the first l_0 moment restrictions out of a total of l_j restrictions under alternative j . This means that if $\delta_{0j} = (\delta_0', 0'_{(j \times 1)})'$, then $m_0(\delta_0) = N_{2j}m_j(\delta_{0j})$, where N_{2j} is defined in (57). It follows that

$$n^{1/2}(\tilde{\delta}_{0j} - \delta_j^*) = -N_{1j}(a_0(\tilde{\delta}_0)'W_0a_0(\tilde{\delta}_0))^{-1}a_0(\tilde{\delta}_0)'W_0N_{2j}n^{1/2}m_j(\delta_0^*) + o_p(1). \quad (61)$$

Inserting (61) into (58) then gives

$$n^{1/2}Dq_j(\tilde{\delta}_{0j}) = T_jn^{1/2}m_j(\delta_j^*) + o_p(1), \quad (62)$$

where T_j is defined in (55).

A central limit theorem implies $n^{1/2}m_j(\delta_j^*) \rightarrow^d N(0_{(h+j \times 1)}, \Omega_j)$, where Ω_j is defined by $\Omega_j = E(f_j(v_i, \delta_j^*)f_j(v_i, \delta_j^*)')$ and f_j is defined in Section 3. It follows that the asymptotic variance of $Dq_j(\tilde{\delta}_{0j})$ can be estimated by $T_j\Omega_jT_j'$. Replacing Ω_j with the estimator B_j defined in (39) yields (54).

A.2 The case of $l_j = h + j$

This section shows that the LM statistic (36) with variance estimator (54) simplifies to (42) in the case where $l_j = h + j$, A_j is invertible and its upper left submatrix is A_0 , and W_j is any nonsingular matrix.

When A_j and W_j are invertible, the LM statistic simplifies to

$$\begin{aligned} R_j &= M_j' W_j A_j (T_j B_j T_j')^{-1} A_j' W_j M_j \\ &= M_j' (U_j B_j U_j')^{-1} M_j, \end{aligned} \quad (63)$$

where

$$U_j = I_{(l_j)} - A_j N_{1j} A_0^{-1} N_{2j}. \quad (64)$$

Partition A_j and A_j^{-1} as

$$A_j = \begin{bmatrix} A_{00} & A_{0j} \\ A_{j0} & A_{jj} \end{bmatrix} \quad \text{and} \quad A_j^{-1} = \begin{bmatrix} A^{00} & A^{0j} \\ A^{j0} & A^{jj} \end{bmatrix}, \quad (65)$$

where A_{00} and A^{00} are h -dimensional and A_{jj} and A^{jj} are j -dimensional square matrices. Assumption 1 implies that $A_{00} = A_0$. (Generally $A^{00} \neq A_0^{-1}$.) Using this result, U_j can be written

$$U_j = \begin{bmatrix} 0_{(h \times h)} & 0_{(h \times j)} \\ -A_{j0} A_{00}^{-1} & I_{(j)} \end{bmatrix}. \quad (66)$$

Partition B_j similarly to A_j . Then

$$U_j B_j U_j' = \begin{bmatrix} 0_{(h \times h)} & 0_{(h \times j)} \\ 0_{(j \times h)} & K_j \end{bmatrix}, \quad (67)$$

where K_j is defined as

$$K_j = A_{j0} A_{00}^{-1} B_{00} (A_{00}^{-1})' A_{j0}' - B_{j0} (A_{00}^{-1})' A_{j0}' - A_{j0} A_{00}^{-1} B_{j0}' + B_{jj}. \quad (68)$$

Partition M_j as

$$M_j = \begin{bmatrix} M_0 \\ M_{\star j} \end{bmatrix}, \quad (69)$$

where $M_0 = 0_{(h \times 1)}$ by the definition of the IV estimator and $M_{\star j}$ is a j -vector. Since K_j is nonsingular, then

$$R_j = M'_{\star j} K_j^{-1} M_{\star j}. \quad (70)$$

Rules for inverting partitioned matrices imply that $A_{j0} A_{00}^{-1} = (A^{jj})^{-1} A^{j0}$. Substituting this into (68) and rearranging yield

$$\begin{aligned} K_j &= (A^{jj})^{-1} A^{j0} B_{00} (A^{j0})' (A^{jj})^{-1'} - B_{j0} (A^{j0})' (A^{jj})^{-1'} - (A^{jj})^{-1} A^{j0} B_{0j} + B_{jj} \\ &= (A^{jj})^{-1} \left(A^{j0} B_{00} (A^{j0})' - A^{jj} B_{j0} (A^{j0})' - A^{j0} B_{0j} (A^{jj})' + A^{jj} B_{jj} (A^{jj})' \right) (A^{jj})^{-1'}. \end{aligned} \quad (71)$$

Recall that $H_j = [0_{(j \times h)} \ I_{(j)}]$. From the last line in the previous expression it follows that

$$K_j = (A^{jj})^{-1} H_j A_j^{-1} B_j (A_j^{-1})' H_j' (A^{jj})^{-1'}. \quad (72)$$

Finally, noting that $H_j A_j^{-1} M_j = A^{jj} M_{\star j}$ yields

$$\begin{aligned} R_j &= M'_{\star j} \left((A^{jj})^{-1} H_j A_j^{-1} B_j (A_j^{-1})' H_j' (A^{jj})^{-1'} \right)^{-1} M_{\star j} \\ &= M'_{\star j} (A^{jj})' \left(H_j A_j^{-1} B_j (A_j^{-1})' H_j' \right)^{-1} A^{jj} M_{\star j} \\ &= M'_j (A_j^{-1})' H_j' \left(H_j A_j^{-1} B_j (A_j^{-1})' H_j' \right)^{-1} H_j A_j^{-1} M_j. \end{aligned} \quad (73)$$

The last line is identical to (42).

A.3 The case of $l_j = l_0$

As noted by e.g. Engle (1984, p795), the form of LM statistics simplifies when the same set of moment restrictions is used both under the null and under the alterna-

tive; that is, when $l_0 = l_j$ and $W_0 = W_j$. Define $E_j = A_j'W_jA_j$ and $G_j = I_{(h+j)} - E_j^{-1/2}H_j'(H_jE_j^{-1}H_j')^{-1}H_jE_j^{-1/2}$. Assumption 1 implies that the first h columns of A_j equal A_0 . Using this fact and formulae for inverting partitioned matrices, it can be verified that

$$N_{1j}(A_0'W_0A_0)^{-1}A_0'W_0N_{2j} = E_j^{-1/2}G_jE_j^{-1/2}A_j'W_j. \quad (74)$$

This result can be used to simplify the expression for T_j in (55),

$$\begin{aligned} T_j &= A_j'W_j[I_{(l_j)} - A_jE_j^{-1/2}G_jE_j^{-1/2}A_j'W_j] \\ &= [I_{(h+j)} - A_j'W_jA_jE_j^{-1/2}G_jE_j^{-1/2}]A_j'W_j \\ &= [I_{(h+j)} - E_j^{1/2}G_jE_j^{-1/2}]A_j'W_j \\ &= [I_{(h+j)} - (I_{(h+j)} - H_j'(H_jE_j^{-1}H_j')^{-1}H_jE_j^{-1})]A_j'W_j \\ &= H_j'(H_jE_j^{-1}H_j')^{-1}H_jE_j^{-1}A_j'W_j. \end{aligned} \quad (75)$$

It follows that $T_jB_jT_j'$ simplifies to

$$T_jB_jT_j' = H_j'(H_jE_j^{-1}H_j')^{-1}H_jE_j^{-1}A_j'W_jB_jW_jA_jE_j^{-1}H_j'(H_jE_j^{-1}H_j')^{-1}H_j. \quad (76)$$

Using the definition of a generalized inverse, it is straightforward to verify that the expression $E_j^{-1}H_j'(H_jE_j^{-1}A_j'W_jB_jW_jA_jE_j^{-1}H_j')^{-1}H_jE_j^{-1}$ is a generalized inverse of $T_jB_jT_j'$. The resulting estimator of $\text{Var}(\text{D}q_j(\tilde{\delta}_{0j}))$ is J_j given in (43). The corresponding LM statistic is given in (45).

References

- Aerts, M., G. Claeskens, and J. D. Hart (1999). Testing the fit of a parametric function. *Journal of the American Statistical Association* 94(447), 869–879.
- Aerts, M., G. Claeskens, and J. D. Hart (2000). Testing lack of fit in multiple regression. *Biometrika* 87(2), 405–424.
- Bierens, H. (1990). A consistent moment test of functional form. *Econometrica* 58(6), 1443–1458.

- Blundell, R., A. Duncan, and K. Pendakur (1998). Semiparametric estimation and consumer demand. *Journal of Applied Econometrics* 13, 435–461.
- Donald, S. G., G. W. Imbens, and W. K. Newey (2003). Empirical likelihood estimation and consistent tests with conditional moment restrictions. *Journal of Econometrics* 117, 55–93.
- Engle, R. F. (1984). Wald, likelihood ratio, and Lagrange multiplier tests in econometrics. In Z. Griliches and M. D. Intriligator (Eds.), *Handbook of Econometrics*, Volume II, pp. 775–826. Amsterdam: Elsevier Science Publishers BV.
- Gallant, A. R. (1981). On the bias in flexible functional forms and an essentially unbiased form: The Fourier flexible form. *Journal of Econometrics* 15, 211–245.
- Hart, J. D. (1997). *Nonparametric smoothing and lack-of-fit tests*. New York: Springer Verlag.
- Horowitz, J. L. (2006). Testing a parametric model against a nonparametric alternative with identification through instrumental variables. *Econometrica* 74(2), 521–538.
- Newey, W. K. and D. McFadden (1994). Large sample estimation and hypothesis testing. In R. F. Engle and D. L. McFadden (Eds.), *Handbook of econometrics. Volume 4. Handbooks in Economics, vol. 2*. Amsterdam; London and New York: Elsevier, North-Holland.
- Pagan, A. and A. Ullah (1999). *Nonparametric econometrics*. Cambridge: Cambridge University Press.
- Tripathi, G. and Y. Kitamura (2003). Testing conditional moment restrictions. *Annals of Statistics* 31(6), 2059–2095.
- Whang, Y.-J. (2001). Consistent specification testing for conditional moment restrictions. *Economics Letters* 71, 299–306.
- White, H. (1982). Maximum likelihood estimation of misspecified models. *Econometrica* 50(1), 1–26.

Table 1: Monte Carlo results for simple IV model (nominal size 5%)

ρ	η	HOR	t	DIN	ACH Min		ACH Same		LM Min		LM Same	
					P	F	P	F	P	F	P	F
<i>Null is true</i>												
<i>Null: $y_i = 0.5x_i + u_i$; DGP: $y_i = 0.5x_i + u_i$</i>												
0.8	0.1	5.1	5.2	4.8	5.2	5.2	5.6	5.0	4.1	4.2	4.2	4.2
0.8	0.5	3.0	3.4	4.3	4.1	3.4	3.5	3.1	3.6	5.6	3.7	5.8
0.7	0.1	4.9	5.2	4.5	5.1	5.2	5.4	4.6	4.3	4.1	4.3	4.1
<i>Null: $y_i = 0.5x_i - 0.5x_i^2 + u_i$; DGP: $y_i = 0.5x_i - 0.5x_i^2 + u_i$</i>												
0.8	0.1	5.3	4.0	4.8	5.0	4.3	5.1	4.2	4.4	4.1	4.5	4.2
0.8	0.5	4.6	7.7	5.0	7.5	3.4	3.8	2.0	5.7	5.7	7.2	6.1
0.7	0.1	5.6	3.6	4.3	5.6	4.3	5.4	4.5	4.5	4.2	4.9	4.3
<i>Null is false</i>												
<i>Null: $y_i = 0.5x_i + u_i$; DGP: $y_i = 0.5x_i - 0.5x_i^2 + u_i$</i>												
0.8	0.1	65.8	71.4	44.7	69.2	69.8	71.1	70.0	39.3	39.3	39.7	39.4
0.8	0.5	72.1	82.7	45.9	78.4	78.2	81.0	79.1	49.9	50.3	50.2	50.4
0.7	0.1	42.1	44.4	25.9	42.1	42.7	45.2	46.9	22.8	22.1	22.8	22.3
<i>Null: $y_i = 0.5x_i + u_i$; DGP: $y_i = 0.5x_i - x_i^2 + x_i^3 + u_i$</i>												
0.8	0.1	68.4	67.1	49.8	64.0	62.7	65.1	64.1	40.3	39.0	40.4	39.0
0.8	0.5	66.3	58.0	48.0	56.6	52.6	55.6	54.5	30.7	34.1	32.0	35.4
0.7	0.1	42.4	41.2	26.2	36.2	36.1	38.3	37.9	17.8	17.1	18.4	16.8
<i>Null: $y_i = 0.5x_i + x_i^2 + u_i$; DGP: $y_i = 0.5x_i - x_i^2 + 4x_i^3 + u_i$</i>												
0.8	0.1	89.0	90.0	72.2	86.8	56.8	93.4	74.8	68.3	65.0	69.1	65.8
0.8	0.5	97.2	98.7	68.5	98.0	82.3	97.7	80.4	83.8	78.0	85.3	79.3
0.7	0.1	52.7	59.0	29.8	49.1	18.2	67.1	34.8	27.6	25.9	29.5	27.2

Legend: HOR: test by Horowitz (2006); t : ordinary t test for adding one additional term to the null model; DIN: the IV test by Donald, Imbens, and Newey (2003); ACH Min: implemented as in Section 2.1; ACH Same: implemented as in Section 2.2; LM Min: the r th LM statistic from the ACH Min calculations; LM Same: the r th LM statistic from the ACH Same calculations; P: based on power basis; F: based on Fourier flexible form basis. *Notes:* HOR, t and DIN quoted from Horowitz (2006). There are 500 observations in each sample and 1000 samples per experiment. In the calculations of the GMM-ACH tests, $r = 6$ and all additional terms under the alternative are orthogonalized to reduce multicollinearity. For the last set of experiments, the dgp process is incorrectly stated in Horowitz's article with the term $2x_i^3$ instead of $4x_i^3$.

Table 2: Engel curve estimates

	Share of total expenditures					
	Food	Fuel	Clothes	Alcohol	Transport	Other
<i>Summary statistics for dependent variable</i>						
Mean	.3565	.0910	.1072	.0606	.1324	.2523
<i>Simple linear model, OLS estimates</i>						
$\tilde{\beta}_1$	-.1338*	-.0472*	.0813*	.0198*	.0394*	.0406*
	(.0060)	(.0032)	(.0059)	(.0041)	(.0069)	(.0067)
<i>Linear model with demographics, OLS estimates</i>						
$\tilde{\beta}_1$	-.1384*	-.0474*	.0819*	.0216*	.0411*	.0412*
	(.0060)	(.0032)	(.0059)	(.0042)	(.0069)	(.0068)
$\tilde{\beta}_2$.0338*	.0012	-.0045	-.0129*	-.0130*	-.0047
	(.0048)	(.0026)	(.0047)	(.0033)	(.0055)	(.0054)
<i>Linear model with demographics, IV estimates</i>						
$\tilde{\beta}_1$	-.1412*	-.0274*	.0473*	.0156	.0295*	.0762*
	(.0122)	(.0067)	(.0123)	(.0085)	(.0142)	(.0140)
$\tilde{\beta}_2$.0341*	-.0005	-.0015	-.0124*	-.0119*	-.0077
	(.0048)	(.0026)	(.0049)	(.0034)	(.0056)	(.0055)
<i>GMM-ACH test of the linear model with demographics</i>						
ACH Min	0.719	6.556*	2.145	0.530	14.268*	3.950
ACH Same	1.200	15.594*	1.013	0.531	16.243*	5.033*
<i>Quadratic model with demographics, IV estimates</i>						
$\tilde{\beta}_1$	-.0618	-2.1008*	.9794	-.0855	2.7383*	-1.4708
	(.6782)	(.5065)	(.6938)	(.4740)	(.9295)	(.8135)
$\tilde{\beta}_2$.0336*	.0112*	-.0068	-.0119*	-.0273*	.0011
	(.0063)	(.0047)	(.0064)	(.0044)	(.0086)	(.0075)
$\tilde{\beta}_3 (w_{1i}^2)$	-.0086	.2256*	-.1014	.0110	-.2947*	.1683
	(.0736)	(.0549)	(.0752)	(.0514)	(.1008)	(.0882)

Legend: $\tilde{\beta}_1$: coefficient on log total expenditure; $\tilde{\beta}_2$: coefficient on indicator of two children; $\tilde{\beta}_3$: coefficient on the square of log total expenditure; ACH Min: implemented using the minimum number of moment restrictions; ACH Same: implemented using the same set of moment restrictions; standard errors in () parentheses; *: statistical significance at the 5% level. *Notes:* Constant included in all models, but not reported. GMM-ACH tests based on a power basis, $r = 6$, and all additional terms under the alternative are orthogonalized to reduce multicollinearity. Data from Blundell, Duncan, and Pendakur (1998). Number of observations: 1519.

- 2009-41: Dennis Kristensen: Pseudo-Maximum Likelihood Estimation in Two Classes of Semiparametric Diffusion Models
- 2009-42: Ole Eiler Barndorff-Nielsen and Robert Stelzer: The multivariate supOU stochastic volatility model
- 2009-43: Lasse Bork, Hans Dewachter and Romain Houssa: Identification of Macroeconomic Factors in Large Panels
- 2009-44: Dennis Kristensen: Semiparametric Modelling and Estimation: A Selective Overview
- 2009-45: Kim Christensen, Silja Kinnebrock and Mark Podolskij: Pre-averaging estimators of the ex-post covariance matrix
- 2009-46: Matias D. Cattaneo, Richard K. Crump and Michael Jansson: Robust Data-Driven Inference for Density-Weighted Average Derivatives
- 2009-47: Mark Podolskij and Mathias Vetter: Understanding limit theorems for semimartingales: a short survey
- 2009-48: Isabel Casas and Irene Gijbels: Unstable volatility functions: the break preserving local linear estimator
- 2009-49: Torben G. Andersen and Viktor Todorov: Realized Volatility and Multipower Variation
- 2009-50: Robinson Kruse, Michael Frömmel, Lukas Menkhoff and Philipp Sibbertsen: What do we know about real exchange rate non-linearities?
- 2009-51: Tue Gørgens, Christopher L. Skeels and Allan H. Würtz: Efficient Estimation of Non-Linear Dynamic Panel Data Models with Application to Smooth Transition Models
- 2009-52: Torben G. Andersen, Dobrislav Dobrev and Ernst Schaumburg: Jump-Robust Volatility Estimation using Nearest Neighbor Truncation
- 2009-53: Florian Heinen, Philipp Sibbertsen and Robinson Kruse: Forecasting long memory time series under a break in persistence
- 2009-54: Tue Gørgens and Allan Würtz: Testing a parametric function against a nonparametric alternative in IV and GMM settings