



SCHOOL OF ECONOMICS AND MANAGEMENT  
FACULTY OF SOCIAL SCIENCES  
AARHUS UNIVERSITY



**CREATES**  
Center for Research in Econometric  
Analysis of Time Series

## CREATES Research Paper 2010-22

### Quantitative Breuer-Major Theorems

Ivan Nourdin, Giovanni Peccati and Mark Podolskij

School of Economics and Management  
Aarhus University  
Bartholins Allé 10, Building 1322, DK-8000 Aarhus C  
Denmark

# Quantitative Breuer-Major Theorems <sup>\*</sup>

Ivan Nourdin<sup>†</sup>  
Paris VI

Giovanni Peccati<sup>‡</sup>  
University of Luxembourg

Mark Podolskij<sup>§</sup>  
ETH Zürich and CREATES

May 12, 2010

## Abstract

We consider sequences of random variables of the type  $S_n = n^{-1/2} \sum_{k=1}^n f(X_k)$ ,  $n \geq 1$ , where  $X = (X_k)_{k \in \mathbb{Z}}$  is a  $d$ -dimensional Gaussian process and  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is a measurable function. It is known that, under certain conditions on  $f$  and the covariance function  $r$  of  $X$ ,  $S_n$  converges in distribution to a normal variable  $S$ . In the present paper we derive several explicit upper bounds for quantities of the type  $|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]|$ , where  $h$  is a sufficiently smooth test function. Our methods are based on Malliavin calculus, on interpolation techniques and on the Stein's method for normal approximation. The bounds deduced in our paper depend only on  $\mathbb{E}[f^2(X_1)]$  and on simple infinite series involving the components of  $r$ . In particular, our results generalize and refine some classic CLTs by Breuer-Major, Giraitis-Surgailis and Arcones, concerning the normal approximation of partial sums associated with Gaussian-subordinated time-series.

**Keywords:** Berry-Esseen bounds; Breuer-Major central limit theorems; Gaussian processes; Interpolation; Malliavin calculus; Stein's method.

**JEL classification:** C60, C10.

---

<sup>\*</sup>Mark Podolskij gratefully acknowledges financial support from CREATES funded by the Danish National Research Foundation.

<sup>†</sup>Laboratoire de Probabilités et Modèles Aléatoires, Université Pierre et Marie Curie, Boîte courrier 188, 4 Place Jussieu, 75252 Paris Cedex 5, France, Email: ivan.nourdin@upmc.fr.

<sup>‡</sup>Faculté des Sciences, de la Technologie et de la Communication; UR en Mathématiques. 6, rue Richard Coudenhove-Kalergi, L-1359 Luxembourg, Email: giovanni.peccati@gmail.com.

<sup>§</sup>Department of Mathematics, ETH Zürich, HG G32.2, 8092 Zürich, Switzerland, Email: mark.podolskij@math.ethz.ch.

# 1 Introduction

Fix  $d \geq 1$  and consider a  $d$ -dimensional centered stationary Gaussian process  $X = (X_k)_{k \in \mathbb{Z}}$ ,  $X_k = (X_k^{(1)}, \dots, X_k^{(d)})$ , defined on a probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ . For any  $1 \leq i, l \leq d$  and  $j \in \mathbb{Z}$ , we denote by

$$r^{(i,l)}(j) = \mathbb{E}[X_1^{(i)} X_{1+j}^{(l)}], \quad (1.1)$$

the covariance of  $X_1^{(i)}$  and  $X_{1+j}^{(l)}$ . Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a measurable function and write

$$S_n = \frac{1}{\sqrt{n}} \sum_{k=1}^n f(X_k), \quad n \geq 1, \quad (1.2)$$

to indicate the sequence of *normalized partial sums* associated with the subordinated process  $k \mapsto f(X_k)$ . One crucial problem in Gaussian analysis is the following:

**Problem P.** *Find conditions on  $f$  and on the covariance  $r$  in order to have that, as  $n \rightarrow \infty$ ,  $S_n$  converges in distribution to a Gaussian random variable.*

Albeit easily stated, Problem P is indeed quite subtle. For instance, as observed e.g. in [8, p. 429], it is in general not possible to deduce a solution to Problem P by using standard central limit results for dependent random variables (for instance, by applying techniques based on mixing). More to the point, slight variations in the form of  $f$  and  $r$  may imply that either the normalization by the factor  $n^{1/2}$  is inappropriate, or the limiting distribution is not-Gaussian (or both): see Dobrushin and Major [16], Rosenblatt [33] and Taqqu [38, 39] for several classic results connected to this phenomenon, as well as Breton and Nourdin [7] for recent developments.

It turns out that an elegant solution to Problem P can be deduced by using the notion of *Hermite rank*. Let  $G \sim N_d(0, I_d)$  and recall that the function  $f$  is said to have Hermite rank equal to  $q \geq 1$ , where  $q$  is an integer, if (a)  $\mathbb{E}[f(G)p_m(G)] = 0$  for every polynomial  $p_m$  (on  $\mathbb{R}^d$ ) of degree  $m \leq q-1$ ; and (b) there exists a polynomial  $p_q$  of degree  $q$  such that  $\mathbb{E}[f(G)p_q(G)] \neq 0$  (see also Proposition 2.1 below). Then, one has the following well-known statement:

**Theorem 1.1 (Breuer-Major theorem for stationary vectors)** *Let  $\mathbb{E}[f^2(X_1)] < \infty$ , and assume that the function  $f$  has Hermite rank equal to  $q \geq 1$ . Suppose that*

$$\sum_{j \in \mathbb{Z}} |r^{(i,l)}(j)|^q < \infty, \quad \forall i, l \in \{1, \dots, d\}. \quad (1.3)$$

*Then  $\sigma^2 := \mathbb{E}[f^2(X_1)] + 2 \sum_{k=1}^{\infty} \mathbb{E}[f(X_1)f(X_{1+k})]$  is well-defined, and belongs to  $[0, \infty)$ . Moreover, one has that*

$$S_n \xrightarrow{d} S \sim N(0, \sigma^2), \quad (1.4)$$

*where  $N(0, \sigma^2)$  indicates a centered Gaussian distribution with variance  $\sigma^2$ , and  $\xrightarrow{d}$  stands for convergence in distribution.*

In the case  $d = 1$ , Theorem 1.1 was first proved by Breuer and Major in [8], whereas Theorem 4 in Arcones [2] proves the statement for a general  $d$  (both proofs in [2, 8] are based on the method of cumulants and diagram formulae – see e.g. [31, 36]). The reader is referred to Sun [35] for an early statement in the case of a Hermite rank equal to 2, and to Giraitis and Surgailis [18] for some continuous-time analogous of Theorem 1.1. Note that any central limit result involving Hermite ranks and series of covariance coefficients is customarily called a ‘Breuer-Major Theorem’, in honor of the seminal paper [8].

Theorem 1.1 and its variations have served as fundamental tools for Gaussian approximations in an impressive number of applications, of which we provide a representative (recent) sample: renormalization of fractional diffusions [1], power variations of Gaussian and Gaussian-related continuous-time processes [3, 4, 15, 21], Gaussian fluctuations of heat-type equations [5], estimation of Hurst parameters of fractional processes [9, 13, 14], unit root problems in econometrics [10], empirical processes of long-memory time-series [23], level functionals of stationary Gaussian fields [20], variations of multifractal random walks [22], and stochastic programming [40]. See also Surgailis [36] for a survey of some earlier uses of Breuer-Major criteria.

Despite this variety of applications, until recently the only available techniques for proving results such as Theorem 1.1 were those based on combinatorial cumulants/diagrams computations. These techniques are quite effective and flexible (see e.g. [24, 34] for further instances of their applicability), but suffer of a fundamental drawback, namely they *do not* allow to deduce Berry-Esseen relations of the type

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| \leq \varphi(n), \quad n \geq 1, \quad (1.5)$$

where  $h$  is a suitable test function, and  $\varphi(n) \rightarrow 0$  as  $n \rightarrow \infty$ . An upper bound such as (1.5) quantifies the error one makes when replacing  $S_n$  with  $S$  for a fixed  $n$ .

In [26, Section 4], the first two authors of this paper proved that one can combine Malliavin calculus (see e.g. [30]) and Stein’s method (see e.g. [12]) to obtain relations such as (1.5) (for some explicit  $\varphi(n)$ ) in the case where: (i)  $d = 1$ , (ii)  $f = H_q$  is a Hermite polynomial of degree  $q \geq 2$  (and thus has Hermite rank equal to  $q$ ), (iii)  $X$  is obtained from the increments of a fractional Brownian motion of Hurst index  $H < 1 - (2q)^{-1}$ , and (iv)  $h$  is either an indicator of a Borel set or a Lipschitz function. Since under (iii) one has that  $|r(j)| \sim j^{2H-2}$ , these findings allow to recover a very special case of Theorem 1.1 (see Example 2.6 below for more details on this point).

The aim of the present work is to extend the techniques initiated in [26] in order to deduce several complete *quantitative Breuer-Major theorems*, that is, statements providing explicit upper bounds such as (1.5) for any choice of  $f$  and  $r$  satisfying the assumptions of Theorem 1.1. We stress by now that we will *not* require that the functions  $f$  enjoy any additional smoothness property, so that our results represent a genuine extension of the findings by Breuer-Major and Arcones.

As anticipated, our techniques are based on the use of Malliavin operators on a Gaussian space, that we combine both with Stein’s method and with an interpolation technique

(already applied in [28, 32]) which is reminiscent of the ‘smart path method’ in Spin Glasses – see e.g. Talagrand [37]. In particular, the use of Stein’s method allows to deal with functions  $h$  that are either Lipschitz or indicators of the type  $h = \mathbf{1}_{(-\infty, z]}$ , whereas the use of interpolations requires test functions that are twice differentiable and with bounded derivatives. Note that this implies that the convergence (1.4) takes indeed place in the stronger topologies of the Kolmogorov and Wasserstein distances.

The rest of the paper is organized as follows. Section 2 contains the statements of our main results, some examples and applications. Section 3 presents some notions and results that are needed to prove our main Theorem 2.2. Section 4 is devoted to proofs.

## 2 Statement of the main results

We keep the assumptions and notation of the previous section. For the sake of notational simplicity, in the following we shall assume that  $X_1 \sim N_d(0, I_d)$ , implying that  $X_k^{(1)}, \dots, X_k^{(d)}$  are independent  $N(0, 1)$  random variables for all  $k \in \mathbb{Z}$ . Note that this is not a restrictive assumption: indeed, by reduction of variables and at the cost of possibly decreasing the value of  $d$ , we may always assume that  $X_k \sim N_d(0, \Sigma)$  for an invertible matrix  $\Sigma \in \mathbb{R}^{d \times d}$ , and by a linear transformation we can further restrict ourselves to the case  $X_k \sim N_d(0, I_d)$ . Observe that, if  $f(x)$  has Hermite rank  $q$  with respect to  $X_1$ , then  $f(Lx)$  has the same Hermite rank  $q$  with respect to  $L^{-1}X_1$ , for every invertible matrix  $L$  – see also [2, p. 2249].

Now, let  $\Lambda$  denote the set of all vectors  $\alpha = (\alpha_1, \dots, \alpha_d)$  with  $\alpha_i \in \mathbb{N} \cup \{0\}$ . For any multi-index  $\alpha \in \Lambda$ , we introduce the notation  $|\alpha| = \sum_{i=1}^d \alpha_i$  and  $\alpha! = \prod_{i=1}^d \alpha_i!$ . When  $\mathbb{E}[f^2(X_1)] < \infty$ , the function  $f$  possesses the unique *Hermite expansion*

$$f(x) = \sum_{\alpha \in \Lambda} a_\alpha \prod_{i=1}^d H_{\alpha_i}(x_i), \quad a_\alpha = (\alpha!)^{-1} \mathbb{E} \left[ f(X_1) \prod_{i=1}^d H_{\alpha_i}(X_1^{(i)}) \right], \quad (2.6)$$

where  $(H_j)_{j \geq 0}$  is the sequence of Hermite polynomials, recursively defined as:  $H_0 = 1$ , and

$$H_j = \delta H_{j-1}, \quad j \geq 1,$$

where  $\delta f(x) = xf(x) - f'(x)$  (for instance:  $H_1(x) = x$ ,  $H_2(x) = x^2 - 1$ , and so on).

The following well-known statement provides a further characterization of Hermite ranks.

**Proposition 2.1** *Let the notation and assumptions of this section prevail (in particular,  $X_1 \sim N_d(0, I_d)$ ). Then, the function  $f$  has Hermite rank  $q \geq 1$  if and only if  $a_\alpha = 0$  for all  $\alpha \in \Lambda$  with  $|\alpha| < q$  and  $a_\alpha \neq 0$  for some  $\alpha \in \Lambda$  with  $|\alpha| = q$ . In particular, if  $f$  has Hermite rank  $q$ , then its Hermite expansion has the form*

$$f(x) = \sum_{m=q}^{\infty} f_m(x), \quad f_m(x) = \sum_{\alpha \in \Lambda: |\alpha|=m} a_\alpha \prod_{i=1}^d H_{\alpha_i}(x_i). \quad (2.7)$$

**Remark on notation.** Fix a function  $f$  such that  $\mathbb{E}[f(X_1^2)] < \infty$  and  $f$  has Hermite rank  $q \geq 1$ . Our main results are expressed in terms of the following collection of coefficients (2.8)–(2.12):

$$\theta(j) = \max_{1 \leq i, l \leq d} |r^{(i, l)}(j)| \quad (2.8)$$

$$K = \inf_{k \in \mathbb{N}} \{\theta(j) \leq d^{-1}, \forall |j| \geq k\} \quad (\text{with } \inf \emptyset = \infty), \quad (2.9)$$

$$\theta = \sum_{j \in \mathbb{Z}} \theta(j)^q, \quad (2.10)$$

$$\sigma_m^2 = \mathbb{E}[f_m^2(X_1)] + 2 \sum_{k=1}^{\infty} \mathbb{E}[f_m(X_1)f_m(X_{1+k})] \quad (\text{for } m \geq q), \quad (2.11)$$

$$\gamma_{n, m, e} = \sqrt{2\theta n^{-1} \sum_{|j| \leq n} \theta(j)^e \sum_{|j| \leq n} \theta(j)^{m-e}} \quad (\text{for } m \geq q \text{ and } 1 \leq e \leq m-1). \quad (2.12)$$

The coefficients  $\theta(j)$ ,  $K$ ,  $\theta$ ,  $\sigma_m^2$  and  $\gamma_{n, m, e}$  will be also combined into the following expressions (2.13)–(2.17):

$$A_{1, n} = \frac{\mathbb{E}[f^2(X_1)]}{2} \left[ \frac{2K^2}{n} + d^q \left( \sum_{|j| \leq n} \theta(j)^q \frac{|j|}{n} + \sum_{|j| > n} \theta(j)^q \right) \right] \quad (2.13)$$

$$A_{2, N} = 2(2K + d^q \theta) \sqrt{\mathbb{E}[f^2(X_1)] \sum_{m=N+1}^{\infty} \mathbb{E}[f_m^2(X_1)]} \quad (2.14)$$

$$A_{3, n, N} = \frac{\mathbb{E}[f^2(X_1)]}{2} \sum_{m=q}^N \left( \frac{d^m}{mm!} \sum_{l=1}^{m-1} ll! \binom{m}{l}^2 \sqrt{(2m-2l)!} \gamma_{n, m, l} \right) \quad (2.15)$$

$$A_{4, n, N} = \frac{\mathbb{E}[f^2(X_1)](2K + d^q \theta)^{1/2}}{2} \times \sum_{q \leq p < s \leq N} d^{s/2} \sqrt{\frac{p!}{s!}} \frac{p+s}{p} \binom{s-1}{p-1} \sqrt{(s-p)!} \gamma_{n, s, s-p}^{1/2} \quad (2.16)$$

$$A_{5, n, N} = \frac{\mathbb{E}[f^2(X_1)]}{2\sqrt{2}} \sum_{q \leq p < s \leq N} (p+s) \sum_{l=1}^{p-1} (l-1)! \binom{p-1}{l-1} \binom{s-1}{l-1} \sqrt{(p+s-2l)!} \times \left( \frac{d^s}{s!} \gamma_{n, s, s-l} + \frac{d^p}{p!} \gamma_{n, p, p-l} \right). \quad (2.17)$$

Note that the coefficients  $K$ ,  $\theta$  and  $\sigma_m$  can in general be infinite, and also that, if  $\mathbb{E}[f(X_1)^2] < \infty$ , if  $f$  has Hermite rank  $q$ , and if (1.3) is in order, then

$$\sigma^2 = \sum_{m=q}^{\infty} \sigma_m^2 < \infty, \quad (2.18)$$

where  $\sigma^2$  is defined in Theorem 1.1.

The next statement, which is the main result of the paper, asserts that the quantities defined above can be used to write explicit bounds of the type (1.5).

**Theorem 2.2 (Quantitative Breuer-Major Theorem)** *Assume that the conditions of Theorem 1.1 are satisfied. Then, the coefficients appearing in formulae (2.8)–(2.17) are all finite. Moreover, the following three bounds are in order.*

(1) *For any function  $h \in C^2(\mathbb{R})$  (that is,  $h$  is twice continuously differentiable) with bounded second derivative, and for every  $n > K$ ,*

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| \leq \|h''\|_{\infty} \left( A_{1,n} + \inf_{N \geq q} \{A_{2,N} + A_{3,n,N} + A_{4,n,N} + A_{5,n,N}\} \right). \quad (2.19)$$

(2) *For any Lipschitz function  $h$ , and for every  $n > K$ ,*

$$\begin{aligned} |\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| &\leq \frac{\|h'\|_{\infty}}{2} \times \\ &\times \left\{ A_{2,n} \left[ \frac{1}{\sigma} + \frac{1}{\sqrt{(2K + d^q \theta) \mathbb{E}[f^2(X_1)]}} \right] + 4 \inf_{N \geq q} \frac{A_{1,n} + A_{3,n,N} + A_{4,n,N} + A_{5,n,N}}{\sqrt{\sum_{m=q}^N \sigma_m^2}} \right\}. \end{aligned} \quad (2.20)$$

(3) *For any  $z \in \mathbb{R}$ , and for every  $n > K$ ,*

$$\begin{aligned} |\mathbb{P}(S_n \leq z) - \mathbb{P}(S \leq z)| &\leq \\ &\frac{\sqrt{2}}{\sigma} \sqrt{A_{2,n} \left[ \frac{1}{\sigma} + \frac{1}{\sqrt{(2K + d^q \theta) \mathbb{E}[f^2(X_1)]}} \right] + 4 \inf_{N \geq q} \frac{A_{1,n} + A_{3,n,N} + A_{4,n,N} + A_{5,n,N}}{\sqrt{\sum_{m=q}^N \sigma_m^2}}}. \end{aligned} \quad (2.21)$$

We will now demonstrate that Theorem 2.2 implies a stronger version of Theorem 1.1, namely that the convergence (1.4) takes place with respect to topologies that are stronger than the one of convergence in distribution. To prove this claim, we need to show in particular that, under the assumptions of Theorem 1.1,  $\gamma_{n,m,e} \rightarrow 0$  as  $n \rightarrow \infty$  for any choice of  $m \geq q$  and  $1 \leq e \leq m - 1$ . This is a consequence of the next Lemma 2.3. In what follows, given positive sequences  $b_n, c_n$ ,  $n \geq 1$ , we shall write  $b_n \lesssim c_n$  whenever  $b_n/c_n$  is bounded, and  $b_n \sim c_n$  if  $b_n \lesssim c_n$  and  $c_n \lesssim b_n$ .

**Lemma 2.3** *Let  $(a_k)_{k \in \mathbb{Z}}$  be a sequence of positive real numbers such that  $\sum_{k \in \mathbb{Z}} a_k^m < \infty$  for some  $m \in \mathbb{N}$ . If  $1 \leq e \leq m - 1$ , then*

$$n^{-1+\frac{e}{m}} \sum_{|k| \leq n} a_k^e \rightarrow 0.$$

*Proof.* Fix  $\delta \in (0, 1)$ , and decompose the sum as  $\sum_{k=1}^n = \sum_{k=1}^{[n\delta]} + \sum_{k=[n\delta]+1}^n$ . By the Hölder inequality we obtain (recall that  $\sum_{k=1}^{\infty} a_k^m$  is finite)

$$n^{-1+e/m} \sum_{k=1}^{[n\delta]} a_k^e \leq n^{-1+e/m} (n\delta)^{1-e/m} \left( \sum_{k=1}^{\infty} a_k^m \right)^{e/m} \leq c\delta^{1-e/m},$$

where  $c$  is some constant, as well as

$$n^{-1+e/m} \sum_{k=[n\delta]+1}^n a_k^e \leq \left( \sum_{k=[n\delta]+1}^n a_k^m \right)^{e/m}.$$

The first term converges to 0 as  $\delta$  goes to zero (because  $1 \leq e \leq m - 1$ ), and the second also converges to 0 for fixed  $\delta$  and  $n \rightarrow \infty$ . This proves the claim.  $\square$

Now recall that, if  $X, Y$  are two real-valued random variables, then the *Kolmogorov distance* between the law of  $X$  and the law of  $Y$  is given by

$$d_{Kol}(X, Y) = \sup_{z \in \mathbb{R}} |\mathbb{P}(X \leq z) - \mathbb{P}(Y \leq z)|. \quad (2.22)$$

If  $\mathbb{E}|X|, \mathbb{E}|Y| < \infty$ , one can also meaningfully define the *Wasserstein distance*

$$d_W(X, Y) = \sup_{f \in \text{Lip}(1)} |\mathbb{E}[f(X)] - \mathbb{E}[f(Y)]|, \quad (2.23)$$

where  $\text{Lip}(1)$  indicates the collection of all Lipschitz functions with Lipschitz constant  $\leq 1$ . Finally, if  $X, Y$  have finite second moments, for every constant  $C > 0$  one can define the distance

$$d_C(X, Y) = \sup_{f \in \mathcal{D}_C^2} |\mathbb{E}[f(X)] - \mathbb{E}[f(Y)]|, \quad (2.24)$$

where  $\mathcal{D}_C^2$  stands for the class of all twice continuously differentiable functions having a second derivative bounded by  $C$ . Note that the topologies induced by  $d_{Kol}$ ,  $d_W$  and  $d_C$ , on the probability measures on  $\mathbb{R}$ , are strictly stronger than the topology of convergence in distribution (see e.g. [17, Ch. 11]).

The next consequence of Theorem 2.2 provides the announced refinement of Theorem 1.1.



**Corollary 2.4 (Breuer-Major, strong version)** *Let the assumptions of Theorem 1.1 prevail. Then, the convergence (1.4) takes place with respect to the three distances  $d_{Kol}$ ,  $d_W$  and  $d_C$  (for all  $C > 0$ ), namely*

$$\lim_{n \rightarrow \infty} d_{Kol}(S_n, S) = \lim_{n \rightarrow \infty} d_W(S_n, S) = \lim_{n \rightarrow \infty} d_C(S_n, S) = 0.$$

*Proof.* Under the assumptions of Theorem 1.1, one has that  $A_{1,n} \rightarrow 0$  as  $n \rightarrow \infty$  (because  $\theta < \infty$ ,  $\sum_{|j| \leq n} \theta(j)^q \frac{|j|}{n} \rightarrow 0$  as  $n \rightarrow \infty$  by bounded convergence). On the other hand, because of (2.18) and since  $\mathbb{E}[f^2(X_1)] < \infty$ , one has that  $A_{2,N} \rightarrow 0$  as  $N \rightarrow \infty$ . Moreover, since  $\gamma_{n,m,e} \rightarrow 0$  for any  $m \geq q$  and  $1 \leq e \leq m - 1$  (due to Lemma 2.3), one has that  $A_{j,n,N} \rightarrow 0$ ,  $j = 3, 4, 5$ , for any fixed  $N$  as  $n \rightarrow \infty$ . We deduce that  $\inf_{N \geq q} \{A_{2,N} + A_{3,n,N} + A_{4,n,N} + A_{5,n,N}\} \rightarrow 0$  as  $n \rightarrow \infty$ . To conclude the proof, it remains to apply (2.19)–(2.21).  $\square$

Next, we present a simplified version of Theorem 2.2 for  $d = 1$  and  $f = H_q$ , where  $H_q$  is the  $q$ th Hermite polynomial. Notice that in this case  $K = 0$ .

**Corollary 2.5 (Hermite subordination)** *Assume that  $d = 1$ ,  $f = H_q$  and  $\sum_{j \in \mathbb{Z}} |r(j)|^q < \infty$ .*

(1) *For any function  $h \in C^2(\mathbb{R})$  with bounded second derivative it holds that*

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| \leq \|h''\|_\infty (A_{1,n} + A_{3,n}). \quad (2.25)$$

(2) *For any Lipschitz function  $h$  it holds that*

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| \leq \frac{2\|h'\|_\infty}{\sigma} (A_{1,n} + A_{3,n}). \quad (2.26)$$

(3) *For any  $z \in \mathbb{R}$  it holds that*

$$|\mathbb{P}(S_n \leq z)] - \mathbb{P}(S \leq z)| \leq \frac{2\|s'_z\|_\infty}{\sigma} (A_{1,n} + A_{3,n}), \quad (2.27)$$

where  $s_z$  is the solution of the Stein's equation associated with the function  $h(x) = 1_{(-\infty, z]}(x)$ , i.e.  $s_z$  solves the differential equation

$$1_{(-\infty, z]}(x) - \Phi(z) = s'_z(x) - xs_z(x), \quad x \in \mathbb{R},$$

with  $\Phi$  being the distribution function of  $N(0, 1)$ . Furthermore, we have that  $\|s'_z\|_\infty \leq 1$  for all  $z \in \mathbb{R}$ .

In this context, the constants  $A_{1,n}$  and  $A_{3,n}$  are given by

$$A_{1,n} = \frac{q!}{2} \theta \left( \sum_{|j| \leq n} |r(j)|^q \frac{|j|}{n} + \sum_{|j| > n} |r(j)|^q \right),$$

$$A_{3,n} = \frac{1}{2q} \sum_{l=1}^{q-1} l! \binom{q}{l}^2 \sqrt{(2q-2l)!} \gamma_{n,q,l},$$

with  $\gamma_{n,q,l}$  defined by (2.12).

*Proof.* From Theorem 3.1 in [26] and Theorem 3.2 in Section 3.3 we obtain the estimate

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| \leq c_h \mathbb{E}|\sigma^2 - \langle DF, -DL^{-1}F \rangle_{\mathfrak{H}}|,$$

where  $c_h = \frac{\|h''\|_\infty}{2}$  in (1),  $c_h = \frac{\|h'\|_\infty}{\sigma}$  in (2) and  $c_h = \frac{\|s'_\sigma\|_\infty}{\sigma}$  in (3). We readily deduce the assertion since  $\mathbb{E}|\sigma^2 - \langle DF, -DL^{-1}F \rangle_{\mathfrak{H}}| \leq 2(A_{1,n} + A_{3,n})$ , which follows from the proof of Theorem 2.2.  $\square$

We remark that the upper bound in (2.27) of Corollary 2.5 is stronger than the general upper bound obtained in (2.21) of Theorem 2.2.

Next, we apply Corollary 2.5 to some particular classes of covariance functions  $r$ .

**Example 2.6 (Covariance functions with polynomial decay)** Assume that  $d = 1$  and  $f = H_q$  with  $q \geq 2$ , and consider a covariance function  $r$  which is regular varying with parameter  $a < 0$ . That is, for all  $|k| \geq 1$ ,  $|r(k)| = |k|^a l(|k|)$ , where  $l$  is a slowly varying function. Recall that for any regular varying function  $r$  with parameter  $\alpha < 0$ , we have the following discrete version of *Karamata's theorem* (see e.g. [6]):

$$\alpha > -1 \quad : \quad \frac{\sum_{k=1}^n |r(k)|}{n^{\alpha+1} l(n)} \rightarrow 1/(\alpha+1),$$

$$\alpha < -1 \quad : \quad \frac{\sum_{k=n}^\infty |r(k)|}{n^{\alpha+1} l(n)} \rightarrow -1/(\alpha+1),$$

as  $n \rightarrow \infty$ . Assume now that  $a < -\frac{1}{q}$ , which implies that the conditions of Theorem 2.2 are satisfied, and  $ae \neq -1$  for any  $e = 1, \dots, q-1$ . By the afore-mentioned convergence results we immediately deduce the following estimates ( $1 \leq e \leq q-1$ )

$$\sum_{|j| \leq n} |r(j)|^q \frac{|j|}{n} \lesssim n^{-1} + n^{aq+1} l(n),$$

$$\sum_{|j| > n} |r(j)|^q \lesssim n^{aq+1} l(n),$$

$$\gamma_{n,q,e} \lesssim n^{-1/2} + n^{a/2} l(n).$$

Thus, for all three cases of Corollary 2.5 we conclude that

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| \lesssim \begin{cases} n^{-1/2} : & a < -1 \\ n^{a/2}l(n) : & a \in (-1, -\frac{1}{q-1}) \\ n^{\frac{aq+1}{2}}l(n) : & a \in (-\frac{1}{q-1}, -\frac{1}{q}) \end{cases}$$

Clearly, the same estimates hold for  $d(S_n, S)$ , where  $d = d_{Kol}$ ,  $d = d_W$  or  $d = d_C$ .

**Example 2.7 (The fractional Brownian motion case)** Let  $d = 1$ ,  $f = H_q$  with  $q \geq 2$  and consider the fractional Gaussian noise  $X_i = B_i^H - B_{i-1}^H$ , where  $B^H$  is a fractional Brownian motion with parameter  $H \in (0, 1)$ . Recall that  $B^H = (B_t^H)_{t \geq 0}$  is a centered Gaussian process (with stationary increments) with covariance structure given by

$$\mathbb{E}[B_t B_s] = \frac{1}{2}(|t|^{2H} + |s|^{2H} - |t + s|^{2H})$$

It is well-known that the correlation function  $r$  of the fractional Brownian noise has the following form:

$$|r(k)| = |k|^{2H-2}l(|k|), \quad k \geq 1,$$

with  $l(|k|) \rightarrow 2H|2H-1|$  as  $|k| \rightarrow \infty$  when  $H \neq \frac{1}{2}$ , and  $l(|k|) = 0$  for  $|k| \geq 1$  when  $H = \frac{1}{2}$ . As in the previous example we immediately deduce that

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| \lesssim \begin{cases} n^{-1/2} : & a \in (-2, -1] \\ n^{a/2} : & a \in [-1, -\frac{1}{q-1}] \\ n^{\frac{aq+1}{2}} : & a \in [-\frac{1}{q-1}, -\frac{1}{q}) \end{cases}$$

with  $a = 2H - 2$  and the same estimates hold for  $d(S_n, S)$  with  $d = d_{Kol}$ ,  $d = d_W$  or  $d = d_C$ . Let us remark that these upper bounds coincide with those derived in Theorem 4.1 in [26].

We finally remark that the rate  $n^{-1/2}$  for  $a = 2H - 2 \in (-2, -1]$  has been proved to be optimal in [27]. For the other two cases the optimality question is still an open problem.

## 3 Toolbox

### 3.1 Malliavin calculus on a Gaussian space

We shall now provide a short introduction to the tools of Malliavin calculus that are needed in the proof of our main Theorem 2.2. The reader is referred to [30] for any unexplained definition or result. Let  $\mathfrak{H}$  be a real separable Hilbert space. We denote by

$W = \{W(h) : h \in \mathfrak{H}\}$  an *isonormal Gaussian process* over  $\mathfrak{H}$ , that is,  $W$  is a centered Gaussian family indexed by the elements of  $\mathfrak{H}$  and such that, for every  $g_1, g_2 \in \mathfrak{H}$ ,

$$\mathbb{E}[W(g_1)W(g_2)] = \langle g_1, g_2 \rangle_{\mathfrak{H}}. \quad (3.28)$$

In what follows, we shall use the notation  $L^2(W) = L^2(\Omega, \sigma(W), \mathbb{P})$ . For every  $q \geq 1$ , we write  $\mathfrak{H}^{\otimes q}$  to indicate the  $q$ th tensor power of  $\mathfrak{H}$ ; the symbol  $\mathfrak{H}^{\odot q}$  indicates the  $q$ th *symmetric* tensor power of  $\mathfrak{H}$ , equipped with the norm  $\sqrt{q!} \|\cdot\|_{\mathfrak{H}^{\otimes q}}$ . We denote by  $I_q$  the isometry between  $\mathfrak{H}^{\odot q}$  and the  $q$ th Wiener chaos of  $X$ . It is well-known (see again [30, Ch. 1]) that any random variable  $F$  belonging to  $L^2(W)$  admits the *chaotic expansion*:

$$F = \sum_{q=0}^{\infty} I_q(f_q), \quad (3.29)$$

where  $I_0(f_0) := E[F]$ , the series converges in  $L^2$  and the kernels  $f_q \in \mathfrak{H}^{\odot q}$ ,  $q \geq 1$ , are uniquely determined by  $F$ . In the particular case where  $\mathfrak{H} = L^2(A, \mathcal{A}, \mu)$ , with  $(A, \mathcal{A})$  a measurable space and  $\mu$  a  $\sigma$ -finite and non-atomic measure, one has that  $\mathfrak{H}^{\odot q} = L^2_s(A^q, \mathcal{A}^{\otimes q}, \mu^{\otimes q})$  is the space of symmetric and square integrable functions on  $A^q$ . Moreover, for every  $f \in \mathfrak{H}^{\odot q}$ ,  $I_q(f)$  coincides with the multiple Wiener-Itô integral (of order  $q$ ) of  $f$  with respect to  $W$  (see [30, Ch. 1]). It is well-known that a random variable of the type  $I_q(f)$ ,  $f \in \mathfrak{H}^{\odot q}$ , has finite moments of all orders (see [19, Ch. VI]). For every  $q \geq 0$ , we write  $J_q$  to indicate the orthogonal projection operator on the  $q$ th Wiener chaos associated with  $W$ , so that, if  $F \in L^2(W)$  is as in (3.29), then  $J_q F = I_q(f_q)$  for every  $q \geq 0$ .

Let  $\{e_k, k \geq 1\}$  be a complete orthonormal system in  $\mathfrak{H}$ . Given  $f \in \mathfrak{H}^{\odot p}$  and  $g \in \mathfrak{H}^{\odot q}$ , for every  $r = 0, \dots, p \wedge q$ , the  $r$ th contraction of  $f$  and  $g$  is the element of  $\mathfrak{H}^{\otimes(p+q-2r)}$  defined as

$$f \otimes_r g = \sum_{i_1, \dots, i_r=1}^{\infty} \langle f, e_{i_1} \otimes \dots \otimes e_{i_r} \rangle_{\mathfrak{H}^{\otimes r}} \otimes \langle g, e_{i_1} \otimes \dots \otimes e_{i_r} \rangle_{\mathfrak{H}^{\otimes r}}. \quad (3.30)$$

In the particular case where  $\mathfrak{H} = L^2(A, \mathcal{A}, \mu)$  (with  $\mu$  non-atomic), one has that

$$f \otimes_r g = \int_{A^r} f(t_1, \dots, t_{p-r}, s_1, \dots, s_r) g(t_{p-r+1}, \dots, t_{p+q-2r}, s_1, \dots, s_r) d\mu(s_1) \dots d\mu(s_r).$$

Moreover,  $f \otimes_0 g = f \otimes g$  equals the tensor product of  $f$  and  $g$  while, for  $p = q$ ,  $f \otimes_p g = \langle f, g \rangle_{\mathfrak{H}^{\otimes p}}$ . Note that, in general, the contraction  $f \otimes_r g$  is *not* a symmetric element of  $\mathfrak{H}^{\otimes(p+q-2r)}$ . The canonical symmetrization of  $f \otimes_r g$  is written  $f \tilde{\otimes}_r g$ . The following multiplication formula is also very useful: if  $f \in \mathfrak{H}^{\odot p}$  and  $g \in \mathfrak{H}^{\odot q}$ , then

$$I_p(f)I_q(g) = \sum_{r=0}^{p \wedge q} r! \binom{p}{r} \binom{q}{r} I_{p+q-2r}(f \tilde{\otimes}_r g). \quad (3.31)$$

Let  $\mathcal{S}$  be the set of all smooth cylindrical random variables of the form

$$F = g(W(\phi_1), \dots, W(\phi_n)),$$

where  $n \geq 1$ ,  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  is a smooth function with compact support and  $\phi_i \in \mathfrak{H}$ . The Malliavin derivative of  $F$  with respect to  $W$  is the element of  $L^2(\Omega, \mathfrak{H})$  defined as

$$DF = \sum_{i=1}^n \frac{\partial g}{\partial x_i}(W(\phi_1), \dots, W(\phi_n)) \phi_i.$$

Also,  $DW(\phi) = \phi$  for every  $\phi \in \mathfrak{H}$ . As usual,  $\mathbb{D}^{1,2}$  denotes the closure of  $\mathcal{S}$  with respect to the norm  $\|\cdot\|_{1,2}$ , defined by the relation

$$\|F\|_{1,2}^2 = \mathbb{E}[F^2] + \mathbb{E}[\|DF\|_{\mathfrak{H}}^2].$$

Note that, if  $F$  is equal to a finite sum of multiple Wiener-Itô integrals, then  $F \in \mathbb{D}^{1,2}$ . The Malliavin derivative  $D$  verifies the following *chain rule*: if  $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}$  is in  $\mathcal{C}_b^1$  (that is, the collection of continuously differentiable functions with bounded partial derivatives) and if  $\{F_i\}_{i=1,\dots,n}$  is a vector of elements of  $\mathbb{D}^{1,2}$ , then  $\varphi(F_1, \dots, F_n) \in \mathbb{D}^{1,2}$  and

$$D\varphi(F_1, \dots, F_n) = \sum_{i=1}^n \frac{\partial \varphi}{\partial x_i}(F_1, \dots, F_n) DF_i.$$

We denote by  $\delta$  the adjoint of the operator  $D$ , also called the *divergence operator*. A random element  $u \in L^2(\Omega, \mathfrak{H})$  belongs to the domain of  $\delta$ , noted  $\text{Dom}\delta$ , if and only if it verifies

$$|\mathbb{E}\langle DF, u \rangle_{\mathfrak{H}}| \leq c_u \|F\|_{L^2} \quad \text{for any } F \in \mathcal{S},$$

where  $c_u$  is a constant depending only on  $u$ . If  $u \in \text{Dom}\delta$ , then the random variable  $\delta(u)$  is defined by the duality relationship (sometimes called ‘integration by parts formula’):

$$\mathbb{E}[F\delta(u)] = \mathbb{E}\langle DF, u \rangle_{\mathfrak{H}}, \quad (3.32)$$

which holds for every  $F \in \mathbb{D}^{1,2}$ .

The operator  $L$ , acting on square integrable random variables of the type (3.29), is defined through the projection operators  $\{J_q\}_{q \geq 0}$  as  $L = \sum_{q=0}^{\infty} -qJ_q$ , and is called the *infinitesimal generator of the Ornstein-Uhlenbeck semigroup*. It verifies the following crucial property: a random variable  $F$  is an element of  $\text{Dom}L$  ( $= \mathbb{D}^{2,2}$ ) if, and only if,  $F \in \text{Dom}\delta D$  (i.e.  $F \in \mathbb{D}^{1,2}$  and  $DF \in \text{Dom}\delta$ ), and in this case:  $\delta DF = -LF$ . Note that a random variable  $F$  as in (3.29) is in  $\mathbb{D}^{1,2}$  if and only if

$$\sum_{q=1}^{\infty} (q+1)! \|f_q\|_{\mathfrak{H}^{\otimes q}}^2 < \infty,$$

and also  $\mathbb{E}[\|DF\|_{\mathfrak{H}}^2] = \sum_{q \geq 1} qq! \|f_q\|_{\mathfrak{H}^{\otimes q}}^2$ . If  $\mathfrak{H} = L^2(A, \mathcal{A}, \mu)$  (with  $\mu$  non-atomic), then the derivative of a random variable  $F$  as in (3.29) can be identified with the element of  $L^2(A \times \Omega)$  given by

$$D_a F = \sum_{q=1}^{\infty} q I_{q-1}(f_q(\cdot, a)), \quad a \in A. \quad (3.33)$$

We also define the operator  $L^{-1}$ , which is the *pseudo-inverse* of  $L$ , as follows: for every  $F \in L^2(W)$ , we set  $L^{-1}F = \sum_{q \geq 1} \frac{1}{q} J_q(F)$ . Note that  $L^{-1}$  is an operator with values in  $\mathbb{D}^{2,2}$  and that  $LL^{-1}F = F - \mathbb{E}[F]$  for all  $F \in L^2(W)$ .

### 3.2 Assessing norms and scalar products

The following statement plays a crucial role in the proof of Theorem 2.2.

**Lemma 3.1** *Let  $F = I_p(h)$  and  $G = I_s(g)$  with  $h \in \mathfrak{H}^{\odot p}$ ,  $g \in \mathfrak{H}^{\odot s}$  and  $p < s$  ( $p, s \geq 1$ ). Then*

$$\text{Var} \left[ \frac{1}{s} \|DG\|_{\mathfrak{H}}^2 \right] = \frac{1}{s^2} \sum_{l=1}^{s-1} l^2 l!^2 \binom{s}{l}^4 (2s-2l)! \|g \tilde{\otimes}_l g\|_{\mathfrak{H}^{\otimes 2s-2l}}^2, \quad (3.34)$$

and

$$\begin{aligned} \mathbb{E} \left[ \left( \frac{1}{s} \langle DF, DG \rangle_{\mathfrak{H}} \right)^2 \right] &\leq p! \binom{s-1}{p-1}^2 (s-p)! \mathbb{E}[F^2] \|g \otimes_{s-p} g\|_{\mathfrak{H}^{\otimes 2p}} \\ &+ \frac{p^2}{2} \sum_{l=1}^{p-1} (l-1)!^2 \binom{p-1}{l-1}^2 \binom{s-1}{l-1}^2 (p+s-2l)! \left( \|h \otimes_{p-l} h\|_{\mathfrak{H}^{\otimes 2l}}^2 + \|g \otimes_{s-l} g\|_{\mathfrak{H}^{\otimes 2l}}^2 \right). \end{aligned} \quad (3.35)$$

*Proof.* [Proof of (3.34)] We have  $DG = sI_{s-1}(g)$  so that, by using (3.31)

$$\begin{aligned} \frac{1}{s} \|DG\|_{\mathfrak{H}}^2 &= s \|I_{s-1}(g)\|_{\mathfrak{H}}^2 = s \sum_{l=0}^{s-1} l! \binom{s-1}{l}^2 I_{2s-2-2l}(g \tilde{\otimes}_{l+1} g) \\ &= s \sum_{l=1}^s (l-1)! \binom{s-1}{l-1}^2 I_{2s-2l}(g \tilde{\otimes}_l g) \\ &= s! \|g\|_{\mathfrak{H}^{\otimes s}}^2 + s \sum_{l=1}^{s-1} (l-1)! \binom{s-1}{l-1}^2 I_{2s-2l}(g \tilde{\otimes}_l g) \\ &= \mathbb{E}[G^2] + s \sum_{l=1}^{s-1} (l-1)! \binom{s-1}{l-1}^2 I_{2s-2l}(g \tilde{\otimes}_l g). \end{aligned}$$

The orthogonality property of multiple integrals leads to (3.34).

[Proof of (3.35)] Thanks once again to (3.31), we can write

$$\begin{aligned}
\langle DF, DG \rangle_{\mathfrak{H}} &= p s \langle I_{p-1}(h), I_{s-1}(g) \rangle_{\mathfrak{H}} \\
&= p s \sum_{l=0}^{p \wedge s-1} l! \binom{p-1}{l} \binom{s-1}{l} I_{p+s-2-2l}(h \tilde{\otimes}_{l+1} g) \\
&= p s \sum_{l=1}^{p \wedge s} (l-1)! \binom{p-1}{l-1} \binom{s-1}{l-1} I_{p+s-2l}(h \tilde{\otimes}_l g).
\end{aligned}$$

It follows that

$$\mathbb{E} \left[ \left( \frac{1}{s} \langle DF, DG \rangle_{\mathfrak{H}} \right)^2 \right] = p^2 \sum_{l=1}^p (l-1)!^2 \binom{p-1}{l-1}^2 \binom{s-1}{l-1}^2 (p+s-2l)! \|h \tilde{\otimes}_l g\|_{\mathfrak{H}^{\otimes(p+s-2l)}}^2. \quad (3.36)$$

If  $l < p$ , then

$$\begin{aligned}
\|h \tilde{\otimes}_l g\|_{\mathfrak{H}^{\otimes(p+s-2l)}}^2 &\leq \|h \otimes_l g\|_{\mathfrak{H}^{\otimes(p+s-2l)}}^2 = \langle h \otimes_{p-l} h, g \otimes_{s-l} g \rangle_{\mathfrak{H}^{\otimes 2l}} \\
&\leq \|h \otimes_{p-l} h\|_{\mathfrak{H}^{\otimes 2l}} \|g \otimes_{s-l} g\|_{\mathfrak{H}^{\otimes 2l}} \\
&\leq \frac{1}{2} (\|h \otimes_{p-l} h\|_{\mathfrak{H}^{\otimes 2l}}^2 + \|g \otimes_{s-l} g\|_{\mathfrak{H}^{\otimes 2l}}^2).
\end{aligned}$$

If  $l = p$ , then

$$\|h \tilde{\otimes}_p g\|_{\mathfrak{H}^{\otimes(s-p)}}^2 \leq \|h \otimes_p g\|_{\mathfrak{H}^{\otimes(s-p)}}^2 \leq \|h\|_{\mathfrak{H}^{\otimes p}}^2 \|g \otimes_{s-p} g\|_{\mathfrak{H}^{\otimes 2p}}.$$

By plugging these last expressions into (3.36), we deduce immediately (3.35).  $\square$

### 3.3 Estimates via interpolations and Stein's method

The forthcoming Theorem 3.2 contains two bounds on normal approximations, that are expressed in terms of Malliavin operators. As anticipated, the proof of Point (1) uses an interpolation technique already applied in [28, 32], which is close to the ‘smart path method’ of Spin Glasses [37]. Point (2) uses estimates from [26].

**Theorem 3.2** *Let  $F$  be a centered element of  $\mathbb{D}^{1,2}$  and let  $Z \sim N(0, \sigma^2)$ ,  $\sigma > 0$ .*

(1) *Suppose that  $h : \mathbb{R} \rightarrow \mathbb{R}$  is twice continuously differentiable and has a bounded second derivative. Then,*

$$|\mathbb{E}[h(F)] - \mathbb{E}[h(Z)]| \leq \frac{\|h''\|_{\infty}}{2} \mathbb{E}|\sigma^2 - \langle DF, -DL^{-1}F \rangle_{\mathfrak{H}}|. \quad (3.37)$$

(2) Suppose that  $h : \mathbb{R} \rightarrow \mathbb{R}$  is Lipschitz. Then,

$$|\mathbb{E}[h(F)] - \mathbb{E}[h(Z)]| \leq \frac{\|h'\|_\infty}{\sigma} \mathbb{E}|\sigma^2 - \langle DF, -DL^{-1}F \rangle_{\mathfrak{H}}|. \quad (3.38)$$

*Proof.* (1) Without loss of generality, we may assume that  $F$  and  $Z$  are independent and defined on the same probability space. Fix  $h$  as in the statement, and define the function  $\Psi(t) = \mathbb{E}[h(\sqrt{1-t}F + \sqrt{t}Z)]$ ,  $t \in [0, 1]$ . Standard results imply that  $\Psi$  is differentiable for every  $t \in (0, 1)$ , and that

$$\Psi'(t) = \frac{1}{2\sqrt{t}} \mathbb{E}[h'(\sqrt{1-t}F + \sqrt{t}Z)Z] - \frac{1}{2\sqrt{1-t}} \mathbb{E}[h'(\sqrt{1-t}F + \sqrt{t}Z)F].$$

By using independence and integration by parts, we obtain immediately that

$$\frac{1}{2\sqrt{t}} \mathbb{E}[h'(\sqrt{1-t}F + \sqrt{t}Z)Z] = \frac{\sigma^2}{2} \mathbb{E}[h''(\sqrt{1-t}F + \sqrt{t}Z)].$$

On the other hand, the relation  $F = LL^{-1}F = -\delta DL^{-1}F$  and (3.32) imply that

$$\begin{aligned} \frac{1}{2\sqrt{1-t}} \mathbb{E}[h'(\sqrt{1-t}F + \sqrt{t}S)F] &= \frac{1}{2\sqrt{1-t}} \mathbb{E}[h'(\sqrt{1-t}F + \sqrt{t}Z)\delta(-DL^{-1}F)] \\ &= \frac{1}{2} \mathbb{E}[h''(\sqrt{1-t}F + \sqrt{t}Z)\langle DF, -DL^{-1}F \rangle_{\mathfrak{H}}]. \end{aligned}$$

The conclusion follows from the fact that

$$|\mathbb{E}[h(F)] - \mathbb{E}[h(Z)]| = |\Psi(1) - \Psi(0)| \leq \int_0^1 |\Psi'(t)| dt.$$

(2) Here we follow the arguments contained in the proof of Theorem 3.1 in [26]. Define  $h_\sigma(x) = h(\sigma x)$ ,  $F_\sigma = \sigma^{-1}F$ , and  $Z_\sigma = \sigma^{-1}Z \sim N(0, 1)$ . Let  $s$  be the solution of the Stein's equation associated with  $h_\sigma$ , i.e.  $s$  solves the differential equation

$$h_\sigma(x) - \mathbb{E}[h_\sigma(Z_\sigma)] = s'(x) - xs(x), \quad x \in \mathbb{R}.$$

It is well-known that such a solution is given by  $s(x) = \frac{1}{\varphi(x)} \int_{-\infty}^x (h_\sigma(t) - \Phi(t))\varphi(t)dt$ , where  $\varphi$  and  $\Phi$  are the density and the distribution function of  $N(0, 1)$ , respectively, and  $\|s'\|_\infty \leq \|h'_\sigma\|_\infty$ . Since  $F_\sigma$  is a centered element of  $\mathbb{D}^{1,2}$  it holds that  $F_\sigma = LL^{-1}F_\sigma = -\delta DL^{-1}F_\sigma$ . By integration by parts formula (3.32) we deduce that

$$\begin{aligned} |\mathbb{E}[h(F)] - \mathbb{E}[h(Z)]| &= |\mathbb{E}[h_\sigma(F_\sigma)] - \mathbb{E}[h_\sigma(Z_\sigma)]| \\ &= |\mathbb{E}[s'(F_\sigma) - F_\sigma s(F_\sigma)]| \\ &= |\mathbb{E}[s'(F_\sigma)(1 - \langle DF_\sigma, -DL^{-1}F_\sigma \rangle_{\mathfrak{H}})]| \\ &\leq \|s'\|_\infty \mathbb{E}|1 - \langle DF_\sigma, -DL^{-1}F_\sigma \rangle_{\mathfrak{H}}| \\ &\leq \|h'_\sigma\|_\infty \mathbb{E}|1 - \langle DF_\sigma, -DL^{-1}F_\sigma \rangle_{\mathfrak{H}}|. \end{aligned}$$



We conclude by using the relations  $\|h'_\sigma\|_\infty = \sigma\|h'\|_\infty$  and

$$\langle DF_\sigma, -DL^{-1}F_\sigma \rangle_{\mathfrak{H}} = \sigma^{-2} \langle DF, -DL^{-1}F \rangle_{\mathfrak{H}}.$$

□

When applied to the special case of a Gaussian random variable  $F$ , Theorem 3.2 yields the following neat estimates.

**Corollary 3.3** *Let  $F \sim N(0, \gamma^2)$  and  $Z \sim N(0, \sigma^2)$ ,  $\gamma, \sigma > 0$ .*

(1) *For every  $h$  twice continuously differentiable and with a bounded second derivative,*

$$|\mathbb{E}[h(F)] - \mathbb{E}[h(Z)]| \leq \frac{\|h''\|_\infty}{2} |\sigma^2 - \gamma^2|. \quad (3.39)$$

(2) *For every Lipschitz function  $h$ ,*

$$|\mathbb{E}[h(F)] - \mathbb{E}[h(Z)]| \leq \frac{\|h'\|_\infty}{\sigma \vee \gamma} |\sigma^2 - \gamma^2|. \quad (3.40)$$

## 4 Proof of Theorem 2.2

### 4.1 Preparation

First, let us remark that the process  $X = (X_k)_{k \in \mathbb{Z}}$  can always be regarded as a subset of an isonormal Gaussian process  $\{W(u) : u \in \mathfrak{H}\}$ , where  $\mathfrak{H}$  is a separable Hilbert space with scalar product  $\langle \cdot, \cdot \rangle_{\mathfrak{H}}$ . More precisely, we shall assume (without loss of generality) that, for every  $k \in \mathbb{Z}$  and every  $1 \leq l \leq d$ , there exists  $u_{k,l} \in \mathfrak{H}$  such that

$$X_k^{(l)} = W(u_{k,l}), \quad \text{and consequently} \quad \langle u_{k,l}, u_{k',l'} \rangle_{\mathfrak{H}} = r^{(l,l')}(k - k'),$$

for every  $k, k' \in \mathbb{Z}$  and every  $1 \leq l, l' \leq d$ . Observe also that  $\mathfrak{H}$  can be taken of the form  $\mathfrak{H} = L^2(A, \mathcal{A}, \mu)$ , where  $\mu$  is  $\sigma$ -finite and non-atomic.

Using the Hermite expansion (2.6) of the function  $f$  we obtain the Wiener chaos representation

$$S_n = \sum_{m=q}^{\infty} I_m(g_m^n), \quad g_m^n \in \mathfrak{H}^{\odot m},$$

where the kernels  $g_m^n$  have the form

$$g_m^n = \frac{1}{\sqrt{n}} \sum_{k=1}^n \sum_{t \in \{1, \dots, d\}^m} b_t u_{k,t_1} \otimes \cdots \otimes u_{k,t_m} \quad (4.41)$$

for certain coefficients  $b_t$  such that the mapping  $t \mapsto b_t$  is symmetric on  $\{1, \dots, d\}^m$ . One also has the identities

$$\mathbb{E}[f_m^2(X_1)] = m! \sum_{t \in \{1, \dots, d\}^m} b_t^2, \quad m \geq q, \quad \mathbb{E}[f^2(X_1)] = \sum_{m=q}^{\infty} m! \sum_{t \in \{1, \dots, d\}^m} b_t^2.$$

Here is a useful preliminary result.

**Lemma 4.1** *For the kernels  $g_m^n$  defined in (4.41) and any  $1 \leq e \leq m - 1$  we obtain the inequality, valid for every  $n$ ,*

$$\|g_m^n \otimes_e g_m^n\|_{\mathfrak{H}^{\otimes 2(m-e)}} \leq \frac{d^m}{m!} \mathbb{E}[f_m^2(X_1)] \gamma_{n,m,e}, \quad (4.42)$$

where  $\gamma_{n,m,e}$  is defined by (2.12). Furthermore, we have that, for every  $n$ ,

$$m! \|g_m^n\|_{\mathfrak{H}^{\otimes m}}^2 \leq \mathbb{E}[f_m^2(X_1)] (2K + d^q \theta), \quad (4.43)$$

where the constants  $K$  and  $\theta$  are defined, respectively in (2.9) and (2.10).

*Proof.* [Proof of (4.42)] Fix  $1 \leq e \leq m - 1$ . Observe that

$$\begin{aligned} g_m^n \otimes_e g_m^n &= \frac{1}{n} \sum_{k_1, k_2=1}^n \sum_{t, s \in \{1, \dots, d\}^m} b_t b_s \prod_{j=1}^e r^{(t_j, s_j)}(k_1 - k_2) \\ &\times u_{k_1, t_{p+1}} \otimes \cdots \otimes u_{k_1, t_m} \otimes u_{k_2, s_{p+1}} \otimes \cdots \otimes u_{k_2, s_m}. \end{aligned}$$

We obtain

$$\begin{aligned} \|g_m^n \otimes_e g_m^n\|_{\mathfrak{H}^{\otimes 2(m-e)}}^2 &\leq \left( \frac{d^m}{m!} \mathbb{E}[f_m^2(X_1)] \right)^2 \\ &\times n^{-2} \sum_{k_1, \dots, k_4=1}^n \theta(k_1 - k_2)^e \theta(k_3 - k_4)^e \theta(k_1 - k_3)^{m-e} \theta(k_2 - k_4)^{m-e}, \end{aligned}$$

where  $\theta(j)$  is defined in (2.8). Since  $\theta(k_3 - k_4)^e \theta(k_1 - k_3)^{m-e} \leq \theta(k_3 - k_4)^m + \theta(k_1 - k_3)^m$  we deduce that

$$\begin{aligned} &n^{-2} \sum_{k_1, \dots, k_4=1}^n \theta(k_1 - k_2)^e \theta(k_3 - k_4)^e \theta(k_1 - k_3)^{m-e} \theta(k_2 - k_4)^{m-e} \\ &\leq 2n^{-1} \sum_{k \in \mathbb{Z}} \theta(k)^m \sum_{|k| \leq n} \theta(k)^e \sum_{|k| \leq n} \theta(k)^{m-e} \leq \gamma_{n,m,e}^2. \end{aligned}$$

Hence, we obtain (4.42).

[Proof of (4.43)] By the Cauchy-Schwarz inequality we have

$$m! \left| \left\langle \sum_{t \in \{1, \dots, d\}^m} b_t u_{k, t_1} \otimes \cdots \otimes u_{k, t_m}, \sum_{t \in \{1, \dots, d\}^m} b_t u_{k+l, t_1} \otimes \cdots \otimes u_{k+l, t_m} \right\rangle_{\mathfrak{H}^{\otimes m}} \right| \leq \mathbb{E}[f_m^2(X_1)]$$

We deduce, for any  $m \geq q$ ,

$$\begin{aligned} m! \|g_m^n\|_{\mathfrak{H}^{\otimes m}}^2 &= \frac{m!}{n} \sum_{k_1, k_2=1}^n \sum_{t, s \in \{1, \dots, d\}^m} b_t b_s \prod_{j=1}^m r^{(t_j, s_j)}(k_1 - k_2) \\ &\leq \left( 2H \mathbb{E}[f_m^2(X_1)] + m! \sum_{|k| \geq K} \theta(k)^m \left( \sum_{t \in \{1, \dots, d\}^m} |b_t| \right)^2 \right) \\ &\leq \mathbb{E}[f_m^2(X_1)] \left( 2K + \sum_{|k| \geq K} (d\theta(k))^m \right) \leq \mathbb{E}[f_m^2(X_1)] (2K + d^q \theta), \end{aligned}$$

which implies (4.43). □

The proofs of Point 1 and Point 2 in Theorem 2.2 are similar, and are detailed in the subsequent two sections.

## 4.2 Proof of Theorem 2.2-(1)

First of all, we remark that  $\theta(j) \rightarrow 0$  as  $|j| \rightarrow \infty$ , because  $\sum_{j \in \mathbb{Z}} \theta(j)^q < \infty$ . This implies that  $K < \infty$ , where the constant  $K$  is defined in (2.9). Moreover, the asymptotic variance  $\sigma^2$  is finite. Indeed we have that  $\sigma^2 = \sum_{m=q}^{\infty} m! \|g_m^n\|_{\mathfrak{H}^{\otimes m}}^2 \leq \mathbb{E}[f^2(X_1)] (2K + d^q \theta)$  due to (4.43).

The main proof is composed of several steps.

**(a) Reduction to a finite chaos expansion.** We start by approximating  $S_n$  by a finite sum of multiple integrals. Define

$$S_{n, N} = \sum_{m=q}^N I_m(g_m^n).$$

Now, let  $h \in C^2(\mathbb{R})$  be a function with bounded second derivative. Since

$$|h(x) - h(y) - h'(0)(x - y)| \leq \frac{1}{2} \|h''\|_{\infty} (y - x)^2 + \|h''\|_{\infty} |x| |y - x|$$

for all  $x, y \in \mathbb{R}$ , we immediately obtain that

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S_{n,N})]| \leq \|h''\|_\infty \left( \frac{1}{2} \|S_n - S_{n,N}\|_{L^2(\mathbb{P})}^2 + \|S_n\|_{L^2(\mathbb{P})} \|S_n - S_{n,N}\|_{L^2(\mathbb{P})} \right).$$

By inequality (4.43) we deduce that

$$\|S_n\|_{L^2(\mathbb{P})}^2 \leq (2K + d^q \theta) \|f(X_1)\|_{L^2(\mathbb{P})}^2$$

and

$$\begin{aligned} \|S_n - S_{n,N}\|_{L^2(\mathbb{P})}^2 &\leq (2K + d^q \theta) \sum_{m=N+1}^{\infty} \|f_m(X_1)\|_{L^2(\mathbb{P})}^2 \\ &\leq (2K + d^q \theta) \|f(X_1)\|_{L^2(\mathbb{P})} \left( \sum_{m=N+1}^{\infty} \|f_m(X_1)\|_{L^2(\mathbb{P})}^2 \right)^{1/2}, \end{aligned}$$

where  $\theta$  is defined by (2.10). We conclude that

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S_{n,N})]| \leq \frac{3(2K + d^q \theta)}{2} \|h''\|_\infty \|f(X_1)\|_{L^2(\mathbb{P})} \left( \sum_{m=N+1}^{\infty} \|f_m(X_1)\|_{L^2(\mathbb{P})}^2 \right)^{1/2}, \quad (4.44)$$

which completes the first step.  $\square$

**(b) Bounds based on the interpolation inequality (3.37).** Let  $Z_N$  be a centered Gaussian random variable with variance  $\sum_{m=q}^N \sigma_m^2$ . By using (3.37) in the special case  $F = S_{n,N}$ ,  $Z = Z_N$  and by applying e.g. (3.33), we obtain

$$\begin{aligned} &\mathbb{E} \left[ h(Z_N) \right] - \mathbb{E} \left[ h(S_{n,N}) \right] \\ &\leq \frac{1}{2} \|h''\|_\infty \left\| \sum_{m=q}^N \sigma_m^2 - \langle DS_{n,N}, -DL^{-1}S_{n,N} \rangle_{\mathfrak{H}} \right\|_{L^2(\mathbb{P})} \\ &\leq \frac{1}{2} \|h''\|_\infty \sum_{p,s=q}^N \left\| \delta_{ps} \sigma_p^2 - s^{-1} \langle DI_p(g_p^n), DI_s(g_s^n) \rangle_{\mathfrak{H}} \right\|_{L^2(\mathbb{P})}, \end{aligned} \quad (4.45)$$

where  $\delta_{ps}$  is the Kronecker symbol. This completes the second step.  $\square$

**(c) The final estimates.** Here we give the approximation of the term on the right-hand side of (4.45). By (4.43) and the dominating convergence theorem we immediately deduce that

$$\mathbb{E}[s^{-1} \|DI_s(g_s^n)\|_{\mathfrak{H}}^2] = s! \|g_s^n\|_{\mathfrak{H}^{\otimes s}}^2 \rightarrow \sigma_s^2 = s! \sum_{k \in \mathbb{Z}} \sum_{t, l \in \{1, \dots, d\}^s} b_t b_l \prod_{j=1}^m r^{(t_j, l_j)}(k).$$

As in the proof of (4.43) we conclude that (recall that we assumed  $n > K$ )

$$\begin{aligned}
|\mathbb{E}[s^{-1}\|DI_s(g_s^n)\|_{\mathfrak{H}}^2] - \sigma_s^2| &\leq s! \sum_{|k|<K} \left| \sum_{t,l \in \{1,\dots,d\}^s} b_t b_l \prod_{j=1}^m r^{(t_j, l_j)}(k) \right| \frac{|k|}{K} \\
&+ s! \sum_{K \leq |k| < n} \left| \sum_{t,l \in \{1,\dots,d\}^s} b_t b_l \prod_{j=1}^m r^{(t_j, l_j)}(k) \right| \frac{|k|}{K} \\
&+ s! \sum_{|k| \geq n} \left| \sum_{t,l \in \{1,\dots,d\}^s} b_t b_l \prod_{j=1}^m r^{(t_j, l_j)}(k) \right| \\
&\leq \mathbb{E}[f_s^2(X_1)] \left\{ \frac{2K^2}{n} + d^q \left( \sum_{|j| \leq n} \theta(j)^q \frac{|j|}{n} + \sum_{|j| > n} \theta(j)^q \right) \right\}.
\end{aligned}$$

Thus we have

$$\begin{aligned}
&\sum_{s=q}^N \left\| \sigma_s^2 - s^{-1}\|DI_s(g_s^n)\|_{\mathfrak{H}}^2 \right\|_{L^2(\mathbb{P})} \\
&\leq \sum_{s=q}^N \left( |\mathbb{E}[s^{-1}\|DI_s(g_s^n)\|_{\mathfrak{H}}^2] - \sigma_s^2| + \sqrt{\text{Var}[s^{-1}\|DI_s(g_s^n)\|_{\mathfrak{H}}^2]} \right) \\
&\leq \mathbb{E}[f_s^2(X_1)] \left\{ \frac{2H^2}{n} + d^q \left( \sum_{|j| \leq n} \theta(j)^q \frac{|j|}{n} + \sum_{|j| > n} \theta(j)^q \right) \right\} \\
&\quad + \sum_{s=q}^N \sqrt{\text{Var}[s^{-1}\|DI_s(g_s^n)\|_{\mathfrak{H}}^2]} \\
&\leq 2(A_{1,n} + A_{3,n,N}),
\end{aligned}$$

where the last inequality follows directly from Lemma 3.1 and 4.1. On the other hand, using the approximation (4.43) and Lemma 3.1 and 4.1 once again we deduce that

$$\begin{aligned}
&\sum_{q \leq p < s \leq N} \left\| s^{-1} \langle DI_p(g_p^n), DI_s(g_s^n) \rangle_{\mathfrak{H}} \right\|_{L^2(\mathbb{P})} + \sum_{q \leq s < p \leq N} \left\| s^{-1} \langle DI_p(g_p^n), DI_s(g_s^n) \rangle_{\mathfrak{H}} \right\|_{L^2(\mathbb{P})} \\
&= \sum_{q \leq p < s \leq N} \frac{p+s}{p} \left\| s^{-1} \langle DI_p(g_p^n), DI_s(g_s^n) \rangle_{\mathfrak{H}} \right\|_{L^2(\mathbb{P})} \leq 2(A_{4,n,N} + A_{5,n,N}).
\end{aligned}$$

Thus, we conclude

$$\left| \mathbb{E} \left[ h \left( \sum_{m=q}^N \sigma_m Y_m \right) - h(S_{n,N}) \right] \right| \leq \|h''\|_{\infty} (A_{1,n} + A_{3,n,N} + A_{4,n,N} + A_{5,n,N}), \quad (4.46)$$

which finishes this step.  $\square$

(d) *Putting things together.* We have the inequality

$$\begin{aligned} |\mathbb{E}[h(S_n)] - \mathbb{E}[h(S)]| &\leq |\mathbb{E}[h(S_n)] - \mathbb{E}[h(S_{n,N})]| + \left| \mathbb{E}[h(S_{n,N})] - \mathbb{E}[h(Z_N)] \right| \\ &\quad + \left| \mathbb{E}[h(Z_N)] - \mathbb{E}[h(S)] \right|. \end{aligned} \quad (4.47)$$

We are left with the derivation of a bound for the third term. By using (3.39) in the special case  $F = Z_N$ , we deduce that

$$\left| \mathbb{E}[h(Z_N)] - \mathbb{E}[h(S)] \right| \leq \frac{1}{2} \|h''\|_\infty \sum_{m=N+1}^{\infty} \sigma_m^2 \leq \frac{2H + d^q \theta}{2} \|h''\|_\infty \sum_{m=N+1}^{\infty} \mathbb{E}[f_m^2(X_1)], \quad (4.48)$$

where the last inequality is deduced by Lemma 4.1. The latter is smaller than  $\frac{1}{4} \|h''\|_\infty A_{2,N}$ , which together with (4.44), (4.45) and (4.46) completes the proof of Theorem 2.2-(1).  $\square$

### 4.3 Proof of Theorem 2.2-(2)

Take  $h$  Lipschitz and write the inequality (4.47). Similar to (4.44) standard computations yield

$$|\mathbb{E}[h(S_n)] - \mathbb{E}[h(S_{n,N})]| \leq \|h'\|_\infty \|S_n - S_{n,N}\|_{L^2(\mathbb{P})} \leq \|h'\|_\infty (2K + d^q \theta)^{1/2} \sqrt{\sum_{m=N+1}^{\infty} \|f_m(X_1)\|_{L^2(\mathbb{P})}^2}.$$

By applying inequality (3.38) in the case  $F = S_{n,N}$ ,  $Z = Z_N$ , one has that

$$\begin{aligned} &\left| \mathbb{E}[h(S_{n,N})] - \mathbb{E}[h(Z_N)] \right| \\ &\leq \frac{\|h'\|_\infty}{\left(\sum_{m=q}^N \sigma_m^2\right)^{1/2}} \sum_{p,s=q}^N \left\| \delta_{ps} \sigma_p^2 - s^{-1} \langle DI_p(g_p^n), DI_s(g_s^n) \rangle_{\mathfrak{H}} \right\|_{L^2(\mathbb{P})} \\ &\leq \frac{\|h'\|_\infty}{\left(\sum_{m=q}^N \sigma_m^2\right)^{1/2}} 2(A_{1,n} + A_{3,n,N} + A_{4,n,N} + A_{5,n,N}), \end{aligned}$$

where the last inequality is obtained by reasoning as in Part (c) of Section 4.2. Finally, an application of inequality (3.40) in the case  $F = Z_N$  yields (since  $\theta \geq 1$ )

$$\left| \mathbb{E}[h(Z_N)] - \mathbb{E}[h(S)] \right| \leq \frac{\|h'\|_\infty}{\sigma} \sum_{m=N+1}^{\infty} \sigma_m^2 \leq \frac{\|h'\|_\infty}{2\sigma} A_{2,N}.$$

Putting the above estimates together yields the desired conclusion.  $\square$

#### 4.4 Proof of Theorem 2.2-(3)

From [12, Theorem 3.1], one can deduce that

$$d_{Kol}(S_n, S) \leq 2\sqrt{d_W(S_n/\sigma, S/\sigma)}.$$

Hence, we get the desired conclusion by combining this inequality with (2.20).  $\square$

## References

- [1] V. V. Anh and N. N. Leonenko (2002): Renormalization and homogenization of fractional diffusion equations with random data. *Probab. Theory Related Fields* **124**(3), 381-408.
- [2] M.A. Arcones (1994): Limit theorems for nonlinear functionals of a stationary Gaussian sequence of vectors. *Ann. Probab.* **22**(4), 2242-2274.
- [3] O.E. Barndorff-Nielsen, J.M. Corcuera and M. Podolskij (2009): Power variation for Gaussian processes with stationary increments. *Stoch. Proc. Appl.* **119**, 1845-1865.
- [4] O.E. Barndorff-Nielsen, J.M. Corcuera and M. Podolskij (2009): Multipower variation for Brownian semi-stationary processes. Preprint.
- [5] L. Beghin, V. P. Knopova, N. N. Leonenko and E. Orsingher (2000): Gaussian limiting behavior of the rescaled solution to the linear Korteweg-de Vries equation with random initial conditions. *J. Statist. Phys.* **99**(3-4) 769-781.
- [6] N.H. Bingham, C.M. Goldie and J.L. Teugels (1987): *Regular variation*. Cambridge University Press.
- [7] J.-C. Breton and I. Nourdin (2008): Error bounds on the non-normal approximation of Hermite power variations of fractional Brownian motion. *Electron. Comm. Probab.* **13**, 482-493 (electronic).
- [8] Breuer, P. and P. Major (1983): Central limit theorems for nonlinear functionals of Gaussian fields. *J. Multivariate Anal.* **13**(3), 425-441.
- [9] C. Berzin and J. León (2007): Estimating the Hurst parameter. *Stat. Inference Stoch. Process.* **10**(1), 49-73.
- [10] B. Buchmann and N.-H. Chan (2009): Integrated functionals of normal and fractional processes. *Ann. Appl. Probab.* **19**(1), 49-70.
- [11] D. Chambers and E. Slud (1989): Central limit theorems for nonlinear functionals of stationary Gaussian processes. *Probab. Th. Rel. Fields* **80**, 323-349.

- [12] L. Chen and Q.-M. Shao (2005): Stein's method for normal approximation. In: *An introduction to Stein's method*, 1-59. Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap. **4**, Singapore Univ. Press, Singapore.
- [13] J.-F. Coeurjolly (2005): Identification of multifractional Brownian motion. *Bernoulli* **11**(6), 987-1008.
- [14] J.-F. Coeurjolly (2008): Hurst exponent estimation of locally self-similar Gaussian processes using sample quantiles. *Ann. Statist.* **36**(3), 1404-1434.
- [15] J. M. Corcuera, D. Nualart and J.H.C. Woerner (2006): Power variation of some integral fractional processes. *Bernoulli* **12**(4), 713-735.
- [16] R.L. Dobrushin and P. Major (1979): Non-central limit theorems for nonlinear functionals of Gaussian fields. *Z. Wahrsch. Verw. Gebiete* **50**(1), 27-52.
- [17] R.M. Dudley (2003): *Real Analysis and Probability* (2<sup>nd</sup> Edition). Cambridge University Press, Cambridge.
- [18] L. Giraitis and D. Surgailis (1985): CLT and other limit theorems for functionals of Gaussian processes. *Z. Wahrsch. verw. Geb.* **70**, 191-212.
- [19] S. Janson (1997): *Gaussian Hilbert Spaces*. Cambridge University Press, Cambridge.
- [20] M. F. Kratz and J. R. León (2001): Central limit theorems for level functionals of stationary Gaussian processes and fields. *J. Theoret. Probab.* **14**(3), 639-672.
- [21] J. León and C. Ludeña (2007): Limits for weighted  $p$ -variations and likewise functionals of fractional diffusions with drift. *Stochastic Process. Appl.* **117**(3), 271-296.
- [22] C. Ludeña (2008):  $L^p$ -variations for multifractal fractional random walks. *Ann. Appl. Probab.* **18**(3), 1138-1163.
- [23] D. Marinucci (2002): The empirical process for bivariate sequences with long memory. *Stat. Inference Stoch. Process.* **8**(2), 205-223.
- [24] D. Marinucci (2007): A Central limit theorem and higher order results for the angular bispectrum. *Probab. Th. Rel. Fields* **141**, 389-409.
- [25] I. Nourdin, D. Nualart and C. A. Tudor (2008): Central and non-central limit theorems for weighted power variations of fractional Brownian motion. *Ann. Inst. H. Poincaré Probab. Statist.*, in press.
- [26] I. Nourdin and G. Peccati (2009): Stein's method on Wiener chaos. *Probab. The. Rel. Fields* **145**(1), 75-118.
- [27] I. Nourdin and G. Peccati (2010): Stein's method and exact Berry-Esseen asymptotics for functionals of Gaussian fields. *Ann. Probab.* **37**(6), 2231-2261.



- [28] I. Nourdin, G. Peccati and G. Reinert (2010): Invariance principles for homogeneous sums: universality of Gaussian Wiener chaos. *Ann. Probab.*, in press.
- [29] I. Nourdin, G. Peccati and A. Réveillac (2008): Multivariate normal approximation using Stein's method and Malliavin calculus. *Ann. Inst. H. Poincaré Probab. Statist.*, in press.
- [30] D. Nualart (2006): *The Malliavin Calculus and Related Topics*. (2nd edition). Springer, Berlin.
- [31] G. Peccati and M.S. Taqqu (2010): *Wiener Chaos: Moments, Cumulants and Diagrams*. Springer, Berlin.
- [32] G. Peccati and C. Zheng (2010): Multi-dimensional Gaussian fluctuations on the Poisson space. Preprint.
- [33] M. Rosenblatt (1961): Independence and dependence. In: *Proc. 4th Berkeley Sympos. Math. Statist. and Prob., Vol. II*, 431-443, Univ. California Press, Berkeley, Calif.
- [34] M. Sodin and B. Tsirelson (2004): Random complex zeroes. I. Asymptotic normality. *Israel J. Math.* **144**, 125-149.
- [35] T.-C. Sun (1965): Some further results on central limit theorems for nonlinear functions of a normal stationary process. *J. Math. Mech.* **14**, 71-85.
- [36] D. Surgailis (2000): CLTs for Polynomials of Linear Sequences: Diagram Formulae with Applications. In: *Long Range Dependence*. Birkhäuser, Basel, 111-128.
- [37] M. Talagrand (2003): *Spin Glasses: a Challenge for Mathematicians. Cavity and Mean fields*. Springer, Berlin.
- [38] M.S. Taqqu (1975): Weak convergence to fractional Brownian motion and to the Rosenblatt process. *Z. Wahrsch. Verw. Gebiete* **31**, 287-302.
- [39] M.S. Taqqu (1979): Convergence of integrated processes of arbitrary Hermite rank. *Z. Wahrsch. Verw. Gebiete* **50**(1), 53-83.
- [40] L. Wang (2002): Asymptotics of statistical estimates in stochastic programming problems with long-range dependent samples. *Math. Methods Oper. Res.* **55**(1), 37-54.

# Research Papers 2010



- 2010-08: Peter R. Hansen and Asger Lunde: Estimating the Persistence and the Autocorrelation Function of a Time Series that is Measured with Error
- 2010-09: Tom Engsted, Thomas Q. Pedersen and Carsten Tanggaard: Pitfalls in VAR based return decompositions: A clarification
- 2010-10: Torben G. Andersen and Luca Benzoni: Stochastic Volatility
- 2010-11: Torben B. Rasmussen: Affine Bond Pricing with a Mixture Distribution for Interest Rate Time-Series Dynamics
- 2010-12: Martin M. Andreasen and Bent Jesper Christensen: The SR Approach: a new Estimation Method for Non-Linear and Non-Gaussian Dynamic Term Structure Models
- 2010-13: Peter Reinhard Hansen, Zhuo (Albert) Huang and Howard Howan Shek: Realized GARCH: Complete Model of Returns and Realized Measures of Volatility
- 2010-14: Bent Jesper Christensen and Michel van der Wel: An Asset Pricing Approach to Testing General Term Structure Models including Heath-Jarrow-Morton Specifications and Affine Subclasses
- 2010-15: Nektarios Aslanidis and Charlotte Christiansen: Smooth Transition Patterns in the Realized Stock Bond Correlation
- 2010-16: Tim Bollerslev and Viktor Todorov: Estimation of Jump Tails
- 2010-17: Ole E. Barndorff-Nielsen, Fred Espen Benth and Almut E. D. Veraart: Ambit processes and stochastic partial differential equations
- 2010-18: Ole E. Barndorff-Nielsen, Fred Espen Benth and Almut E. D. Veraart: Modelling energy spot prices by Lévy semistationary processes
- 2010-19: Jeroen V.K. Rombouts and Lars Stentoft: Multivariate Option Pricing with Time Varying Volatility and Correlations
- 2010-20: Charlotte Christiansen: Intertemporal Risk-Return Trade-off in
- 2010-21: Marco Aiolfi, Carlos Capistrán and Allan Timmermann : Forecast Combinations
- 2010-22: Ivan Nourdin, Giovanni Peccati and Mark Podolskij: Quantitative Breuer-Major Theorems